



Calhoun: The NPS Institutional Archive
DSpace Repository

Theses and Dissertations

1. Thesis and Dissertation Collection, all items

1953

Nuclear magnetic moments of nitrogen(14),
chlorine(37), and indium(115)

Biard, Forrest Rosecrans

Ohio State University

<http://hdl.handle.net/10945/14465>

Downloaded from NPS Archive: Calhoun



<http://www.nps.edu/library>

Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.


Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

Nuclear Magnetic Moments of Nitrogen¹⁴,
Chlorine³⁷, and Indium¹¹⁵

M.S. Thesis

Forrest R. Biard

THESIS
B51



THE HISTORY OF THE
CITY OF BOSTON

FROM THE FIRST SETTLEMENT OF THE COLONY
TO THE PRESENT TIME

BY
JOSEPH NEALE

NEW YORK:
1857.

NUCLEAR MAGNETIC MOMENTS OF NITROGEN¹⁴,
CHLORINE³⁷, AND INDIUM¹¹⁵

Ph.D., Ohio State University, 1953

Department of Physics
(Approved by Faculty Committee)

Abstract of

A Thesis

Presented in Partial Fulfillment of the Requirements
for the Degree Master of Science

By

FORREST ROSECRANS BIARD, B.S.

The Ohio State University
1953

Approved by:

Adviser

THESIS
BSI

Submitted to the Faculty of the
University of Wisconsin

In partial
fulfillment of the requirements for the
degree of

MASTERS

in the Department of
Psychology

by

J. J. GILBERT

MASTERS OF ARTS
1927

1927

1927

NUCLEAR MAGNETIC MOMENTS OF NITROGEN¹⁴,

CHLORINE³⁷, AND INDIUM¹¹⁵

FORREST ROBERTS BIRD

B.S., United States Naval Academy, 1934

Department of Physics
(Approved by Dudley Williams)

The "extra-nuclear" properties of the nucleus are discussed and the importance of accurate measurements of these properties in testing future nuclear theories is pointed out. The various methods used to measure one of these properties, the nuclear magnetic moment, are described.

Nuclear magnetic moments μ_I of Nitrogen¹⁴, Chlorine³⁷, and Indium¹¹⁵ were measured by means of a magnetic resonance spectrometer of the super-regenerative type. Improvements of the modulation methods and of the magnetic field homogeneity of the spectrometer previously employed are discussed.

The values obtained for the various μ_I were

$$\begin{array}{lll} \mu_I(N^{14}) & (0.40361 \pm 0.00002) \mu_n, & \\ \mu_I(Cl^{37}) & (0.683722 \pm 0.000048) \mu_n, & \text{and} \\ \mu_I(In^{115}) & (5.50945 \pm 0.00015) \mu_n. & \end{array}$$

These results are observed to compare satisfactorily with those previously obtained by investigators at Stanford using another type of magnetic resonance spectrometer. Results obtained by these methods in the cases of In¹¹³ and In¹¹⁵ are seen not to agree with those obtained by the atomic beam method. An explanation offered

THE UNIVERSITY OF CHICAGO

DEPARTMENT OF CHEMISTRY

RESEARCH REPORT

NO. 1000

(Submitted by the author)

The following report was prepared by the author and his associates in the Department of Chemistry, University of Chicago, during the period from 1945 to 1947. It is a preliminary report of the results of the work done during this period. The work was supported by the National Science Foundation, Office of Naval Research, and the Army Research Office.

The work was carried out in the Department of Chemistry, University of Chicago, during the period from 1945 to 1947. It is a preliminary report of the results of the work done during this period. The work was supported by the National Science Foundation, Office of Naval Research, and the Army Research Office.

The work was carried out by the author and his associates.

1. The work was carried out by the author and his associates.	2. The work was carried out by the author and his associates.
3. The work was carried out by the author and his associates.	4. The work was carried out by the author and his associates.
5. The work was carried out by the author and his associates.	6. The work was carried out by the author and his associates.

The work was carried out by the author and his associates. It is a preliminary report of the results of the work done during this period. The work was supported by the National Science Foundation, Office of Naval Research, and the Army Research Office.

by Foley for this lack of agreement is summarized.

The results are further compared with those predicted by the Schmidt equations and by using in these same equations the value of the angular momenta of the odd proton for Cl^{37} and In^{115} obtained from nuclear shell theory. A discussion of the relationships between μ_I and the nuclear quadrupole moment Q is also given with the conclusion being reached that to date no theory has yet been successful in establishing any important relationships between these two quantities.

Presented as Partial Fulfillment of the Requirements
for the Degree Master of Science

FRANK J. SCHUBERT, JR.

The Ohio State University

NUCLEAR MAGNETIC MOMENTS OF NITROGEN¹⁴,
CHLORINE³⁷, AND INDIUM¹¹⁵

A Thesis

Presented in Partial Fulfillment of the Requirements
for the Degree Master of Science

By

FORREST ROSECRANS BIARD, B.S.

The Ohio State University
1953

Approved by:

Adviser

REPORT OF THE COMMISSIONER OF THE GENERAL LAND OFFICE

IN RESPONSE TO A RESOLUTION OF THE HOUSE OF COMMONS

1871

Presented in pursuance of an Order of the House of Commons
for the Deputy Master of the Rolls

By

JOHN R. HARRISON, Esq.

THE ONLY LAND SURVEYOR

LONDON

Printed by

TABLE OF CONTENTS

	Page
I. Introduction	1
(a) Nuclear Mass	3
(b) Nuclear Spin	3
(c) Nuclear Magnetic Moments	5
(d) Nuclear Quadrupole Moments	8
II. Methods of Measuring Magnetic Moments	
(a) Hyperfine Structure	11
(b) Molecular Beams	13
(c) Work on Liquids and Solids	16
(1) Method of Purcell and Pound	24
(2) Method of Bloch	29
(3) Super-regenerative Method	32
(4) Heterodyne Method	38
III. Experimental Arrangements	
(a) Apparatus	43
(b) Modulation Improvements	46
(c) Homogenizing of the Magnetic Field	47
IV. Experimental Results	58
(a) Indium ¹¹⁵	59
(b) Chlorine ³⁷	60
(c) Nitrogen ¹⁴	61
V. Discussion of Results	62
(a) Shell Theory	65
(b) Relationships Between μ_1 and Q	69

1	Introduction	1
2	(a) History of the	2
3	(b) History of the	3
4	(c) History of the	4
5	(d) History of the	5
6	(e) History of the	6
7	(f) History of the	7
8	(g) History of the	8
9	(h) History of the	9
10	(i) History of the	10
11	(j) History of the	11
12	(k) History of the	12
13	(l) History of the	13
14	(m) History of the	14
15	(n) History of the	15
16	(o) History of the	16
17	(p) History of the	17
18	(q) History of the	18
19	(r) History of the	19
20	(s) History of the	20
21	(t) History of the	21
22	(u) History of the	22
23	(v) History of the	23
24	(w) History of the	24
25	(x) History of the	25
26	(y) History of the	26
27	(z) History of the	27
28	(aa) History of the	28
29	(ab) History of the	29
30	(ac) History of the	30
31	(ad) History of the	31
32	(ae) History of the	32
33	(af) History of the	33
34	(ag) History of the	34
35	(ah) History of the	35
36	(ai) History of the	36
37	(aj) History of the	37
38	(ak) History of the	38
39	(al) History of the	39
40	(am) History of the	40
41	(an) History of the	41
42	(ao) History of the	42
43	(ap) History of the	43
44	(aq) History of the	44
45	(ar) History of the	45
46	(as) History of the	46
47	(at) History of the	47
48	(au) History of the	48
49	(av) History of the	49
50	(aw) History of the	50
51	(ax) History of the	51
52	(ay) History of the	52
53	(az) History of the	53
54	(ba) History of the	54
55	(bb) History of the	55
56	(bc) History of the	56
57	(bd) History of the	57
58	(be) History of the	58
59	(bf) History of the	59
60	(bg) History of the	60
61	(bh) History of the	61
62	(bi) History of the	62
63	(bj) History of the	63
64	(bk) History of the	64
65	(bl) History of the	65
66	(bm) History of the	66
67	(bn) History of the	67
68	(bo) History of the	68
69	(bp) History of the	69
70	(bq) History of the	70
71	(br) History of the	71
72	(bs) History of the	72
73	(bt) History of the	73
74	(bu) History of the	74
75	(bv) History of the	75
76	(bw) History of the	76
77	(bx) History of the	77
78	(by) History of the	78
79	(bz) History of the	79
80	(ca) History of the	80
81	(cb) History of the	81
82	(cc) History of the	82
83	(cd) History of the	83
84	(ce) History of the	84
85	(cf) History of the	85
86	(cg) History of the	86
87	(ch) History of the	87
88	(ci) History of the	88
89	(cj) History of the	89
90	(ck) History of the	90
91	(cl) History of the	91
92	(cm) History of the	92
93	(cn) History of the	93
94	(co) History of the	94
95	(cp) History of the	95
96	(cq) History of the	96
97	(cr) History of the	97
98	(cs) History of the	98
99	(ct) History of the	99
100	(cu) History of the	100

I. INTRODUCTION

The present status of the theory of the atomic nucleus is roughly comparable to that of the extra-nuclear atom circa 1912, by which time a mass of spectroscopic and x-ray data had been assembled and a number of empirical relationships established, yet a satisfactory model of the electronic configuration had not been proposed. However, within the ensuing few decades this wealth of data sufficed to produce first the Bohr model of the hydrogen atom and later the Lande vector model, the concept of closed electronic shells and the introduction of electron and nuclear spins, all of which contributed greatly to the explanation in classical or semi-classical concepts of a great many of the phenomena observed involving bound electrons. The later of these developments were more or less paralleled by de Broglie's proposal of the wave nature of matter in 1925 and the development of matrix mechanics and wave mechanics by Heisenberg and Schroedinger, respectively, within the following one or two years.

The status of the theory of the nuclear atom is today quite similar to the above situation. The data assembled by a host of experimenters in recent years relative to nuclear energy levels has, particularly with the aid of wave mechanics, enabled physicists to offer a certain number of fairly plausible explanations of basic nuclear phenomena. Even so, many other basic phenomena are not yet adequately explained and no theory yet proposed appears to hold forth very great expectations. One of the new attempts to systematize the prediction of nuclear properties is the fairly recently proposed theory of closed shells in nuclei, but this theory as yet has produced certainly no

The present volume of the history of the science of man is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series.

It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series.

It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series.

It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series. It is a very important one, and it is one of the most important of the series.

more than a limited number of satisfactory predictions of nuclear properties.

One of the great handicaps suffered by the nuclear physicist is the difficulty he encounters in trying to obtain accurate data pertaining to nuclear energy levels. Fairly precise measurement of alpha particle energies is difficult, but really accurate measurement of beta particle, gamma ray, and neutron energies presents even greater problems, whereas the measurement of many of the lower energy phenomena is now, perhaps, impossible. Indeed, in many cases the experimenter is greatly handicapped by not being able to identify accurately the nuclide from which phenomena involving radioactivity originate.

In contrast to this general cloud of less than moderate accuracy which surrounds quantitative measurements of nuclear phenomena, there are certain observable quantities of the nucleus referred to frequently as "extra-nuclear properties" which are capable of being measured with remarkable precision using techniques now available. These are the properties of nuclear charge, nuclear mass, nuclear spin, nuclear magnetic moment and, to a less accurate degree of measurement, the nuclear quadrupole moment. The nuclear charge for those elements occurring naturally on earth has already been completely determined in terms of the electronic charge. It may reasonably be expected that any satisfactory theory of the nucleus must account adequately and accurately for all of these extra-nuclear properties as a first test because of the superior accuracy of data assembled or being assembled relative to these properties.

...the ...

© 1999 Blackwell Science Ltd *Journal of Internal Medicine* 245: 111–117

(a) Nuclear Mass

Absolute measurements of nuclear masses are now limited by the accuracy with which Avogadro's number has been determined. However, relative mass determinations made with the mass spectrograph may be obtained accurate to one part in fifteen thousand in the case of light atoms and to one part in ten thousand for atoms of greater mass number.¹ This accuracy is sufficient to measure mass changes in all

1. Halliday, D. Introductory Nuclear Physics. New York: John Wiley & Sons, 1950, p. 258.

but very low energy nuclear reactions involving emission of γ -rays. By the use of Einstein's formula for the equivalence of mass and energy one can from mass spectrographic data obtain the energies involved in a great number of nuclear reactions.

(b) Nuclear Spin

Nuclear spin was first proposed by Pauli in 1924 to explain hyperfine structures (excluding isotopic effect) by assuming that the nucleus possessed a magnetic moment which perturbed the motion of the orbital electrons.² Not until 1936, however, when Goudsmit and

2. Pauli, W. Naturwissenschaften 12, 741 (1924).

Uhlenbeck gave adequate explanation of the possibilities of accounting for fine structure in atomic spectra by assigning to the electron

ALL OF WHICH ARE THE RESULT OF THE FOLLOWING FACTS:—
First, the fact that the American people are not properly
informed as to the true condition of the country and
the state of the nation.

The second fact is the fact that the American people are not properly
informed as to the true condition of the country and
the state of the nation.

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION
PUBLISHED WEEKLY CHICAGO, ILL., MAY 1, 1914

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION
PUBLISHED WEEKLY CHICAGO, ILL., MAY 1, 1914

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION
PUBLISHED WEEKLY CHICAGO, ILL., MAY 1, 1914

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION
PUBLISHED WEEKLY CHICAGO, ILL., MAY 1, 1914

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION
PUBLISHED WEEKLY CHICAGO, ILL., MAY 1, 1914

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION
PUBLISHED WEEKLY CHICAGO, ILL., MAY 1, 1914

an inherent spin* of $\frac{1}{2} \left(\frac{h}{2\pi} \right)$ did it become evident that nuclear spin might indeed explain certain hyperfine structures. Using the concept of space quantization of the nuclear spin vector together with the vector model of the extra-nuclear atom one could ascertain that the inherent part of the nuclear angular momentum might be determined merely from the number of hyperfine components in a spectral line. This inherent or spin angular momentum of the nucleus is given by $I \left(\frac{h}{2\pi} \right)$, where I is called the "nuclear spin" and has zero, integral or half-integral values. For a given nucleus in the ground state I has a fixed value.*

Another method of determining nuclear spins is afforded by observing the optical spectra of diatomic molecules containing identical nuclei, for in such spectra there is an alternation of intensity in the band spectra in the ratio of $(I+1)/(I)$. Thus the measurement of relative intensity of alternate lines in band spectra of certain diatomic molecules can be used to determine nuclear spin.

The investigation of nuclear spins has revealed that spins of the elementary nuclear particles, neutrons and protons, in the ground state have spin angular momentum values of $\frac{1}{2} \left(\frac{h}{2\pi} \right)$. Furthermore the

*The lengths of the nuclear spin and the electronic spin angular momentum vectors are shown by the wave mechanics to have the magnitudes $\sqrt{I(I+1)}$ and $\sqrt{s(s+1)}$, respectively, in units of $\left(\frac{h}{2\pi} \right)$. However, the maximum projected values they may possess in any specified direction, say, that determined by an external magnetic field, are I and s .

$$\left(\frac{1}{\pi}\right)^{\frac{1}{2}}$$

$$I\left(\frac{1}{\pi}\right)$$

$$\left(\frac{1}{\pi}\right)^{\frac{1}{2}}$$

$$\left(\frac{1}{\pi}\right)^{\frac{1}{2}}$$

Pauli exclusion principle applies in the case of these individual elementary particles of a nucleus as well as for the bound atomic electrons. It has also been determined empirically that a nucleus with an even number of nucleons will have the values $I = 0, 1, 2, \dots$, whereas one with odd mass number will have $I = 1/2, 3/2, 5/2, \dots$. In so far as present day observations have revealed it may also be said that nuclei having an even number of both protons and neutrons have, at least in their ground states, zero resultant angular momenta and zero nuclear magnetic moments.

(c) Nuclear Magnetic Moment

According to classical electromagnetic theory the magnetic moment of a spinning charge cloud with mass m and charge e uniformly distributed throughout its mass should be $\frac{e}{2mc} p_s$, where p_s is the angular momentum due to spin. This classical theory is not borne out by experiment, which reveals a magnetic moment of value $2 \cdot \frac{e}{2mc} p_s$ for the electron due to its spin, which is just twice the classical prediction. The 2 in this last expression is called the g -factor. This expression for the magnetic moment of the spinning electron can be conveniently expressed as

$$\mu_s = 2 \cdot \frac{e}{2mc} \cdot p_s = 2 \cdot \frac{e h}{4\pi mc} \cdot s = 2 \cdot s \cdot \mu_B,$$

in which μ_B , called the Bohr magneton, has the value

$$\mu_B = \frac{e h}{4\pi mc}$$

and s , the spin angular momentum of the electron, has the value $\frac{1}{2}$ expressed in units of $(\frac{h}{2\pi})$. Wave mechanics offers an adequate explanation of magnetic effects associated with the orbital angular momentum

The first part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \frac{1}{2} \log \frac{1+x}{1-x}$. It is shown that $f(x)$ is an odd function and that its Taylor series expansion around $x=0$ is given by $f(x) = x + \frac{1}{3}x^3 + \frac{1}{5}x^5 + \dots$. The second part of the paper is devoted to the study of the function $g(x) = \frac{1}{2} \log \frac{1+x}{1-x}$ and its properties. It is shown that $g(x)$ is an even function and that its Taylor series expansion around $x=0$ is given by $g(x) = \frac{1}{2}x^2 + \frac{1}{8}x^4 + \frac{1}{16}x^6 + \dots$.

3. The function $f(x)$ and its properties

The function $f(x)$ is defined by the equation $f(x) = \frac{1}{2} \log \frac{1+x}{1-x}$. It is shown that $f(x)$ is an odd function and that its Taylor series expansion around $x=0$ is given by $f(x) = x + \frac{1}{3}x^3 + \frac{1}{5}x^5 + \dots$. The function $f(x)$ is also shown to be a solution of the differential equation $(1-x^2)f'(x) = 1$. The function $f(x)$ is also shown to be a solution of the functional equation $f(x) + f\left(\frac{1}{x}\right) = \frac{1}{2} \log \frac{1+x}{1-x} + \frac{1}{2} \log \frac{1+\frac{1}{x}}{1-\frac{1}{x}}$.

$$f(x) = \frac{1}{2} \log \frac{1+x}{1-x} = \frac{1}{2} \log \frac{1+x}{1-x} = \frac{1}{2} \log \frac{1+x}{1-x} = \frac{1}{2} \log \frac{1+x}{1-x}$$

$$f(x) = \frac{1}{2} \log \frac{1+x}{1-x} = \frac{1}{2} \log \frac{1+x}{1-x} = \frac{1}{2} \log \frac{1+x}{1-x} = \frac{1}{2} \log \frac{1+x}{1-x}$$

$$f(x) = \frac{1}{2} \log \frac{1+x}{1-x} = \frac{1}{2} \log \frac{1+x}{1-x} = \frac{1}{2} \log \frac{1+x}{1-x} = \frac{1}{2} \log \frac{1+x}{1-x}$$

The function $f(x)$ is also shown to be a solution of the differential equation $(1-x^2)f'(x) = 1$. The function $f(x)$ is also shown to be a solution of the functional equation $f(x) + f\left(\frac{1}{x}\right) = \frac{1}{2} \log \frac{1+x}{1-x} + \frac{1}{2} \log \frac{1+\frac{1}{x}}{1-\frac{1}{x}}$.

of electrons, but the explanation of the value observed for the magnetic moment of the spinning electron is explained at the present only by Dirac's relativistic wave mechanics, and this explanation now appears to be too small by 12 parts in ten thousand.^{3,4}

3. Kusch, P., and Foley, H.M. Phys. Rev. 72, 1326 (1947).

4. Lamb, H.E., and Retherford, R.C. Phys. Rev. 72, 241 (1947).

By analogy with that of the electron it might appear that the magnetic moment of the proton due to its spin should be $2 \frac{e}{2Mc} \cdot \mu_I$, where M is the mass of the proton and μ_I the spin angular momentum. If this prediction were correct the proton would have a magnetic moment roughly one two-thousandth that of the electron, since both particles have spin $\frac{1}{2} \left(\frac{h}{2\pi} \right)$ in the ground state. Although this analogy is not borne out by experiment, it is nevertheless convenient to measure nuclear magnetic moments in units of

$$\mu_n = \frac{eh}{4\pi Mc}.$$

The unit μ_n is called the nuclear magneton. Since nuclear magnetic moments differ only in magnitude and possibly sign from the prediction given in

$$\mu_I = \frac{e}{2Mc} \cdot \mu_I = I \cdot \frac{eh}{4\pi Mc} = I \mu_n,$$

it is common practice to introduce a nuclear g-factor defined by the equation

$$\mu_I = g_I \cdot \frac{e}{2Mc} \cdot \mu_I = g_I \frac{eh}{4\pi Mc} \cdot I = g_I I \mu_n,$$

where μ_I represents $I \left(\frac{h}{2\pi} \right)$. In general g_I is not the same for any

The following information was obtained from the records of the Department of Health and Human Services, Office of the Assistant Secretary for Health Policy and Statistics, Division of Health Policy and Statistics, Bureau of Health Care Statistics, Office of Health Data Collection and Analysis, Office of Health Data Collection and Analysis, Office of Health Data Collection and Analysis, Office of Health Data Collection and Analysis.

$$x_{\text{M}} = \frac{1}{2}$$

$$\frac{1}{2} = \frac{1}{2}$$

...and even was offering the students in the public schools

$$\text{new } I = \frac{29}{50000} \cdot I = 19 \cdot \frac{1}{50000} = 19$$

© 2000 Blackwell Science Ltd, *Journal of Internal Medicine* 247: 399–406

$$I_{\text{eff}} = I \cdot \frac{A_0}{A_{\text{max}}} = I_0 \cdot \frac{1}{2} = \frac{I_0}{2}$$

$\frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4}$

two different types of nuclei.

It is informative to consider the magnetic moment of the deuteron, the simplest compound nucleus. To do this first consider the individual magnetic moments of its constituents, the neutron and the proton.

Experiment has revealed that the magnetic moment of the proton is $(2.7926 \pm 0.00006) \mu_n$.⁵ Since the spin of the proton is $\frac{1}{2}$, one ob-

5. Sumner, H., Thomas, R.A. and Hipple, J.A. Phys. Rev. 80, 407 (1950).

tains the value 5.58536 for its nuclear g-factor. The neutron, an uncharged particle by classical theory expected to have zero magnetic moment, nevertheless has been observed to have a spin magnetic moment⁶ of $(-1.9128 \pm 0.00009) \mu_n$. Its g-factor is thus seen

6. Zach, J.T. Rev. Mod. Phys. 22, 64 (1950).

to be $(-)5.5856$. In the deuteron the projected spin vectors of the proton and the neutron are parallel and give a value of unity for the nuclear spin. Thus one might suppose that the magnetic moment of the deuteron would be given by the algebraic sum of the moments of the two elementary particles, which would be

$$\mu(D^2) = \mu(P') + \mu(n') = 0.87988 \mu_n.$$

However, experiment gives a magnetic moment⁷ of $0.85738 \mu_n$ for the

7. Aichler, H. A Table of Nuclear Moment Data (Preliminary Issue). 1952, p. 13.

The first part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \sum_{n=0}^{\infty} a_n x^n$. It is shown that $f(x)$ is analytic in the region $|x| < 1$ and that it satisfies the functional equation $f(x) = x f(x^2) + g(x)$, where $g(x)$ is a certain function. The second part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \sum_{n=0}^{\infty} a_n x^n$. It is shown that $f(x)$ is analytic in the region $|x| < 1$ and that it satisfies the functional equation $f(x) = x f(x^2) + g(x)$, where $g(x)$ is a certain function.

The third part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \sum_{n=0}^{\infty} a_n x^n$. It is shown that $f(x)$ is analytic in the region $|x| < 1$ and that it satisfies the functional equation $f(x) = x f(x^2) + g(x)$, where $g(x)$ is a certain function. The fourth part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \sum_{n=0}^{\infty} a_n x^n$. It is shown that $f(x)$ is analytic in the region $|x| < 1$ and that it satisfies the functional equation $f(x) = x f(x^2) + g(x)$, where $g(x)$ is a certain function.

The fifth part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \sum_{n=0}^{\infty} a_n x^n$. It is shown that $f(x)$ is analytic in the region $|x| < 1$ and that it satisfies the functional equation $f(x) = x f(x^2) + g(x)$, where $g(x)$ is a certain function. The sixth part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \sum_{n=0}^{\infty} a_n x^n$. It is shown that $f(x)$ is analytic in the region $|x| < 1$ and that it satisfies the functional equation $f(x) = x f(x^2) + g(x)$, where $g(x)$ is a certain function.

The seventh part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \sum_{n=0}^{\infty} a_n x^n$. It is shown that $f(x)$ is analytic in the region $|x| < 1$ and that it satisfies the functional equation $f(x) = x f(x^2) + g(x)$, where $g(x)$ is a certain function. The eighth part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \sum_{n=0}^{\infty} a_n x^n$. It is shown that $f(x)$ is analytic in the region $|x| < 1$ and that it satisfies the functional equation $f(x) = x f(x^2) + g(x)$, where $g(x)$ is a certain function.

deuteron, a difference of $0.02250 \mu_n$, which may be considered precise to 0.5%. Thus the last equation is not true. Theoretical explanations for this difference have been offered but the problem can by no means be considered completely solved. Nor has any acceptable theoretical justification yet been proposed for the values obtained by experiment for the magnetic moments of the proton and the neutron nor for any compound nucleus when its spin is other than zero. For most nuclei with higher mass numbers and non-zero spin it is usually impossible to predict even approximately the magnitudes of their magnetic moments.

Present techniques for measuring relative magnetic moments of nuclei commonly give an accuracy of five or six significant figures. This accuracy of measurement is rivaled only by those made with the mass spectrograph, so it is hoped that nuclear magnetic moment determinations may in the future prove to be of as great or greater value in the formulation of nuclear theory as relative mass determinations have been in giving to us our present knowledge of nuclear forces.

(d) Nuclear Quadrupole Moment

The nuclear quadrupole moment is a measure of the extent by which the charge of a nucleus varies from spherical symmetry. The expression for the quadrupole moment may be obtained briefly in the following manner.⁸

8. Halliday, D. Introductory Nuclear Physics. New York: John Wiley & Sons, 1950, pp. 59-63.

If a point in space is chosen as an origin and from there a straight line is drawn as a reference axis, the expression for the scalar potential at points along the axis due to a finite number of discrete point charges near the origin can be obtained as a function of R , the distances on the axis from the origin. This scalar potential ϕ can always be expressed by means of the series

$$\phi = c_1 R^{-1} + c_2 R^{-2} + c_3 R^{-3} + c_4 R^{-4} + \dots,$$

in which c_n is the component of the electrical multipole moment of order 2^{n-1} for the particular configuration, origin and direction. In the above series the quadrupole term is therefore seen to be represented by $\frac{c_3}{R^3}$.

In nuclei the discrete charges are protons, and the nuclear center of mass and the axis of nuclear spin are taken as the origin and the reference axis, respectively.

If the discrete system of charges is changed to a continuous charge distribution of uniform density δ with center of charge somewhere near the origin, one obtains as the expression for the scalar potential the series

$$\phi = R^{-1} \iiint \delta d\tau + R^{-2} \iiint z \delta d\tau + R^{-3} \iiint (3z^2 - a^2) \delta d\tau + \dots$$

In this equation " a " is the radial distance of the volume element, dv , from the center of the charge distribution, and z is the component along the spin axis of the radius vector from the origin to the volume element. The magnitudes of the various order multipole terms give indications of the distribution of charge about the particular origin and axis of reference.

It is shown that the system is stable in the sense of Lyapunov.

For values outside of the 1000-100,000 range, the following equation

Copyright © 2004 John Wiley & Sons, Ltd.

Information on the use of the model can be found in the user manual.

© 2004 The Authors
Journal compilation © 2004 Blackwell Publishing Ltd

$$\phi = c_1 R^{-1} + c_2 R^{-2} + c_3 R^{-3} + c_4 R^{-4} + \dots$$

ALL INFORMATION CONTAINED HEREIN IS UNCLASSIFIED

For the above results the following steps in the derivation lead to the results:

216

Downloaded At: 11:53 11 September 2009

—some systems by volume 6619 2 —

Copyright © 2004 by John Wiley & Sons, Inc.

$$\dots + \ln 2 \left(\frac{1}{2} - \omega \right) \left(\frac{1}{2} \right)^{\frac{1}{2}} + \ln 2 \left(\frac{1}{2} \right)^{\frac{1}{2}} + \ln 2 \left(\frac{1}{2} \right)^{\frac{1}{2}} = \phi$$

presently under all 50 states and the District of Columbia. The 1990s will also see the

and the other two are the same as in the first case.

The dipole term will always be zero for any charge distribution possessing mirror symmetry about a plane perpendicular to the axis and passing through the origin. The dipole term is zero for all nuclei.

Nuclei with zero spin have no preferred axis, so the charge distribution is effectively symmetric with the result that such nuclei do not possess quadrupole moments. Wave mechanics can be used to prove that the quadrupole moment is zero also when the nuclear spin has the value one-half.⁹ Hence no nucleus with spin less than unity

9. Blatt, J.M., and Weisskopf, V.F. Theoretical Nuclear Physics. New York: John Wiley & Sons, 1952, pp. 26-30.

can possess a quadrupole moment other than zero. For other nuclei, a potential function with a negative quadrupole term indicates an oblate spheroidal distribution of charge about the axis of spin, whereas a potential function with positive quadrupole term indicates a prolate spheroidal distribution.

In nuclear theory the quadrupole component is defined as

$$Q = \frac{1}{2e} \iiint (3z^2 - a^2) \rho d\tau.$$

Experimental determinations of Q give values¹⁰ ranging from $Q = 7.0 \times 10^{-24} \text{ cm}^2$ for $_{71}\text{Lu}^{176}$ to $(-)1.2 \times 10^{-24} \text{ cm}^2$ for $_{51}\text{Sb}^{72}$.

10. Mack, J.E. Revs. Mod. Phys. 22, 64 (1950).

It has not yet been possible to demonstrate experimentally any conclusive evidence of octupole or higher order electric moments in nuclei.

The first part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \int_0^x f(t) dt$. It is shown that $f(x)$ is a continuous function and that it satisfies the differential equation $f'(x) = f(x)$. The second part of the paper is devoted to the study of the properties of the function $g(x)$ defined by the equation $g(x) = \int_0^x g(t) dt$. It is shown that $g(x)$ is a continuous function and that it satisfies the differential equation $g'(x) = g(x)$.

The third part of the paper is devoted to the study of the properties of the function $h(x)$ defined by the equation $h(x) = \int_0^x h(t) dt$. It is shown that $h(x)$ is a continuous function and that it satisfies the differential equation $h'(x) = h(x)$.

The fourth part of the paper is devoted to the study of the properties of the function $k(x)$ defined by the equation $k(x) = \int_0^x k(t) dt$. It is shown that $k(x)$ is a continuous function and that it satisfies the differential equation $k'(x) = k(x)$.

The fifth part of the paper is devoted to the study of the properties of the function $l(x)$ defined by the equation $l(x) = \int_0^x l(t) dt$. It is shown that $l(x)$ is a continuous function and that it satisfies the differential equation $l'(x) = l(x)$.

$$f(x) = \int_0^x f(t) dt$$

The sixth part of the paper is devoted to the study of the properties of the function $m(x)$ defined by the equation $m(x) = \int_0^x m(t) dt$. It is shown that $m(x)$ is a continuous function and that it satisfies the differential equation $m'(x) = m(x)$.

The seventh part of the paper is devoted to the study of the properties of the function $n(x)$ defined by the equation $n(x) = \int_0^x n(t) dt$. It is shown that $n(x)$ is a continuous function and that it satisfies the differential equation $n'(x) = n(x)$.

The eighth part of the paper is devoted to the study of the properties of the function $o(x)$ defined by the equation $o(x) = \int_0^x o(t) dt$. It is shown that $o(x)$ is a continuous function and that it satisfies the differential equation $o'(x) = o(x)$.

II. METHODS OF MEASURING MAGNETIC MOMENTS

(a) Hyperfine Structure

Approximate values of nuclear magnetic moments may be determined from hyperfine structure in atomic spectra by using a method developed by Goudsmit.¹¹ Since this method involves certain approximations and

11. Goudsmit, S. Phys. Rev. 43, 636 (1933).

some correction factors which are themselves inaccurately known, it is perhaps the least accurate method for this purpose. The method remains of value even today, however, for not all nuclear magnetic moments have yet been determined by more precise methods. Such is the case with the isotopes of silver, for which the only values so far obtained are those calculated from hyperfine structure data.

The hyperfine splitting of an energy level with total electronic angular momentum ($Jh/2\pi$) is given by

$$\delta \nu = A I J \cos(IJ) = \frac{1}{2} A \left\{ F(F+1) - J(J+1) - I(I+1) \right\},$$

in which I is the nuclear angular momentum in units of $(h/2\pi)$, F is the resultant of I and J , and the cosine is a "quantum" cosine. The proportionality factor A is equal to the distance between two adjacent hyperfine levels divided by the largest of their F values and is given a positive sign when the larger F value belongs to the higher energy. A is proportional to the nuclear g -factor.

For a single non-penetrating electron it is customary to write

(2) *Generalization*

The generalization of the results obtained in the preceding section to the case of a general system of linear equations is as follows:

$$A^{-1} = \frac{1}{\Delta} \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

where Δ is the determinant of the matrix A , and a_{ij} are the elements of the adjoint matrix. The elements a_{ij} are defined as the minors of the matrix A with respect to the element a_{ji} . The adjoint matrix is the transpose of the matrix of minors. The elements a_{ij} are defined as the minors of the matrix A with respect to the element a_{ji} . The adjoint matrix is the transpose of the matrix of minors. The elements a_{ij} are defined as the minors of the matrix A with respect to the element a_{ji} . The adjoint matrix is the transpose of the matrix of minors.

$$\Delta = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

The determinant Δ is the sum of the products of the elements of the matrix A with the elements of the adjoint matrix. The determinant Δ is the sum of the products of the elements of the matrix A with the elements of the adjoint matrix. The determinant Δ is the sum of the products of the elements of the matrix A with the elements of the adjoint matrix. The determinant Δ is the sum of the products of the elements of the matrix A with the elements of the adjoint matrix. The determinant Δ is the sum of the products of the elements of the matrix A with the elements of the adjoint matrix.

"a" instead of A and the following relation holds,

$$a = \frac{R g^2 Z^3}{n^3 (l + \frac{1}{2}) j(j+1)} \cdot \frac{g_I}{1838} \text{ cm}^{-1}.$$

For a single outer electron in a penetrating orbit, the expression for "a" becomes

$$a = \frac{R g^2 Z_i Z_o^2}{n_o^3 (l + \frac{1}{2}) j(j+1)} \cdot \frac{g_I}{1838} \text{ cm}^{-1}$$

in which n_o is the principal quantum number, Z_o is the effective nuclear charge when the electron is outside the core electrons, and Z_i the average effective nuclear charge when the electron has penetrated the core.

The ordinary spin doublets for non-s electrons are obtained from the formula

$$\Delta \nu = \frac{R g^2 Z_i^2 Z_o^2}{n_o^3 l(l+1)} \text{ cm}^{-1},$$

which, combined with the previous equation gives

$$a = \frac{\Delta \nu}{Z_i (l + \frac{1}{2})} \cdot \frac{l(l+1)}{j(j+1)} \cdot \frac{g_I}{1838} \text{ cm}^{-1}.$$

After applying certain relativity corrections shown to be necessary by Breit¹² and by Racah¹³ and solving for g_I , one obtains for s-electrons

12. Breit, G. Phys. Rev. 30, 463 (1931).

13. Racah, G. Matte. F. Physik 71, 431 (1931).

$$g_I = \frac{3a}{8Rg^2} \cdot \frac{n_o^3}{Z_i Z_o^2} \cdot \frac{1838}{K(\frac{1}{2}, Z_i)}.$$

For non-s electrons, the result is

$$g_I = \frac{a Z_i}{\Delta \nu} \cdot \frac{j(j+1)(l + \frac{1}{2})}{l(l+1)} \cdot \frac{\lambda(l, Z_i)}{K(j, Z_i)} \cdot 1838.$$

$$\omega = \frac{1}{1838} \cdot \frac{1}{(1+j)\left(\frac{1}{2}+j\right)} = 0$$

$$\omega = \frac{1}{1838} \cdot \frac{1}{(1+j)\left(\frac{1}{2}+j\right)} = 0$$

$$\omega = \frac{1}{1838} \cdot \frac{1}{(1+j)\left(\frac{1}{2}+j\right)} = 0$$

$$\omega = \frac{1}{1838} \cdot \frac{1}{(1+j)\left(\frac{1}{2}+j\right)} = 0$$

$$\omega = \frac{1}{1838} \cdot \frac{1}{(1+j)\left(\frac{1}{2}+j\right)} = 0$$

$$\omega = \frac{1}{1838} \cdot \frac{1}{(1+j)\left(\frac{1}{2}+j\right)} = 0$$

$\kappa(j, z_i)$ is the relativity correction by which the equation for the hyperfine structure must be multiplied, and $\lambda(l, z_i)$ is a similar correction for the multiplet separation. They are determined by the formulas

$$\kappa(j, z_i) = 4j(j + \frac{1}{2})(j+1)/(4\rho^2 - 1)\rho, \quad \rho^2 = (j + \frac{1}{2})^2 - (\alpha z_i)^2,$$

$$\lambda(l, z_i) = [2l(l+1)/(\alpha z_i)^2] \left\{ \left[(l + \frac{1}{2})^2 - (\alpha z_i)^2 \right]^{1/2} - \left[l^2 - (\alpha z_i)^2 \right]^{1/2} \right\}.$$

In many-electron spectra the same electron may be responsible for part or all of the interaction which produces the hyperfine splitting of several different levels. If "a" is calculated for this electron from the various hyperfine structures and these values turn out to be consistent with each other, this agreement of values for "a" is not necessarily sufficient to warrant the assumption that the approximation made in the various formulas above are valid. Only if the values of g_I calculated from different electrons of the same atom agree should one assume that the approximations involved are admissible.

(b) Molecular Beams

In the molecular beam method as developed by Rabi and his co-workers¹⁴ at Columbia University for measuring nuclear magnetic moments,

14. Rabi, I.I., Millman, S., Kusch, P., and Zacharias, J.R.

Phys. Rev. 55, 526 (1939).

a well collimated beam of neutral molecules is passed through the fields

of three magnets, A, C and B, in a highly evacuated space as shown in Figure 1. The source of the molecules is a small oven O which contains the appropriate substance and from which molecules with a modified Maxwellian speed distribution escape through a small slit. These escaping molecules are collimated by additional slits so that all molecules entering the field of magnet A do so within a very small angle to a line parallel to the length of the pole faces. A further defining slit S is located between magnets A and C. At a small distance past the exit from magnet B on the axis defined by the collimating slits is a detector for measuring the intensity of the molecular beam.

The field of magnet A is of the order of 12,000 gauss and is very inhomogeneous (approximately 10^5 gauss/cm.), the magnet field H and the field gradient (dH/dz) in this magnet being in opposite directions. Magnet B is similar to A except that it is arranged to produce a field H in the same direction as, but has its gradient opposite in direction to that of A. Magnet C, which is between A and B, produces a uniform field in the same direction as those of magnets A and B. A hairpin loop to which can be applied a very weak oscillating radio-frequency field at right angles to the magnetic field is installed between the pole faces of magnet C.

In molecular beam experiments one employs molecules in a Σ state in which the resultant electronic angular momentum is zero to the first order. It has been calculated that the interactions between the nuclear magnetic moments in a molecule in a Σ state and between these and the magnetic moment associated with molecular rotation involve




FIGURE 1.
THE PATHS OF FOUR MOLECULES IN A MOLECULAR-BEAM
MAGNETIC RESONANCE APPARATUS

magnetic fields of the order of 100 gauss or less.¹⁴ External magnetic fields of several thousand gauss will decouple the nuclear spins from each other and also from the molecular rotation to the extent that the nuclear spins may be regarded as free.

A nucleus of spin I when in a magnetic field may assume one of only $2I+1$ different directions with respect to the direction of the magnetic field.* In an inhomogeneous magnetic field of the type

*Some of the basic concepts introduced briefly here will be treated in considerable detail in the next section.

shown for magnets A and B a nucleus with magnetic moment experiences a resultant force $\mu_z \frac{dH}{dz}$, where μ_z is the component of μ in the direction of the field H .

Consider now a molecule which has entered the field of magnet A with just the proper velocity, spin orientation and deviation from the dotted axis shown in Figure 1 to be deflected so as to pass through slit S into the homogeneous field of magnet C. If the nuclear spin undergoes no change in spatial quantization while the molecule is passing through the field of C, the arrangement is such that the molecule will be deflected back toward the axis just sufficiently to impinge upon the detector D after emerging from the field of the third magnet.

After operating conditions have been properly adjusted and the beam of molecules impinges on the detector D, a radio frequency oscillation is applied to the hairpin loop in the field of magnet C.

of the new federal financial system, which will be
based on the new federal financial system, which will be

Now, the energy difference ΔE between adjacent levels resulting from the nuclear spin axis changing its projection in the direction of the magnetic field by $\Delta M_I = \pm 1$ is

$$\Delta E = g_I \mu_n H.$$

It therefore appears plausible that the proper application of a radio-frequency field of frequency given by $h\nu = \Delta E$ might produce transitions between adjacent levels. Indeed, transitions involving reorientation of the nuclear spin vector with respect to the magnetic field involving either emission or absorption are induced when the frequency of the r-f oscillations applied to the hairpin loop has the resonance value

$$\nu = \frac{\Delta E}{h} = \frac{g_I \mu_n H}{h}.$$

Thus when the proper field strength H exists for the r-f oscillations to be of the frequency required for resonance, transitions involving reorientation of the direction of nuclear spin axes will occur. Molecules whose nuclei undergo such reorientation of spin axis while passing through the field of magnet C will no longer be deflected by magnet B so as to impinge upon the detector but will be deflected so as to miss it as shown by the dotted trajectories in Figure 1. Thus, when the resonance condition is realized there is a decided decrease in detector "current". The nuclear g-factor of the isotope being studied may then be obtained from the relations

$$\Delta E = h\nu = g_I \mu_n H, \quad \text{and}$$

$$g_I = \frac{h\nu}{\mu_n H}.$$

From the above relations it will be noted that the frequency of radiation required to produce transitions in the field of C is independent of the particular orientation of the nuclear spin axis. For most nuclides the resonance frequency is of the order of several megacycles per second for fields of some 2,000 to 12,000 gauss.

The Rabi method was the first to give values of nuclear magnetic moments to four significant figures. The limitation upon accuracy in this method is the requirement that the strength of the magnetic field be determined. Other disadvantages of the molecular beam method are the requirements for a high vacuum throughout the magnet system and for a fairly high temperature in the oven. But it also has certain advantages, among them being the fact that very minute quantities of an isotope are required for measurements.

This method has largely been replaced by the more accurate nuclear resonance spectrometers, but its usefulness has by no means disappeared.

(c) Work on Liquids and Solids

The next significant advance in the technique of measuring nuclear magnetic moments came in 1946 when Purcell, Pound and Torrey at Harvard¹⁵ and Bloch, Hansen and Packard at Stanford¹⁶ simultane-

15. Purcell, E.M., Torrey, H.G., and Pound, R.V. Phys. Rev. **62**, 37 (1946).

16. Bloch, F., Hansen, W.W., and Packard, Martin. Phys. Rev. **62**, 127 (1946).

from the same situation is still in mind that the Department of
Education reported in various publications in the year 1910 to 1911
regarding the condition of the schools in the State. The
report contains the following statement: "It is found in many
instances that the schools are in a state of neglect and
disrepair. The buildings are old and the equipment is
outdated. The teachers are poorly paid and the
schools are overcrowded. The results of the examination
show that the pupils are not learning as much as they
should. The Department of Education is now making
a study of the situation and will report to the
Legislature in the next session." The Department of
Education is now making a study of the situation and
will report to the Legislature in the next session.

(a) The Department of Education is now making a study of the situation and will report to the Legislature in the next session.

The Department of Education is now making a study of the situation and will report to the Legislature in the next session.

The Department of Education is now making a study of the situation and will report to the Legislature in the next session.

(b) The Department of Education is now making a study of the situation and will report to the Legislature in the next session.

The Department of Education is now making a study of the situation and will report to the Legislature in the next session.

(c) The Department of Education is now making a study of the situation and will report to the Legislature in the next session.

cusly and independently developed new resonance techniques requiring equipment appreciably more simple than that used in molecular beam methods. A distinct advantage of these new methods lies in the fact that it frequently is unnecessary to alter the physical or chemical form of the sample, thus avoiding many of the experimental difficulties of the molecular beam method. In addition to their use in nuclear studies these new methods provide a means for investigating the establishment of the thermal equilibrium essential to magnetic methods for attaining very low temperatures and for obtaining information concerning crystal structure, phase transitions in solids, and hindered internal motions in solids.

The basic semi-classical concept which most readily describes the essential phenomena involved in resonance experiments will now be discussed in detail.

As previously noted, the nuclear magnetic moment may be expressed as

$$\mu_I = g_I \cdot \frac{e}{2Mc} \cdot p_I.$$

Results from wave mechanics indicate that the value of $|p_I|$ is $[I(I+1)]^{\frac{1}{2}} \left(\frac{h}{2\pi}\right)$ but that the maximum projected value which $|p_I|$ may assume in the direction of an applied external magnetic field is

$$(p_H)_{\max} = I \left(\frac{h}{2\pi}\right).$$

The permitted values of p_I , the projected magnitude of p_I upon the direction of the field, are

$$p_I = M_I \left(\frac{h}{2\pi}\right),$$

where $M_I = I, I-1, I-2, \dots, -I+1, -I$.

If a magnet of dipole moment μ is placed in a magnetic field H ,

and independently derived our various theoretical results
 which are especially true when the case is concerned
 with a system of three or more particles. It is true
 that it is necessary to assume in order to obtain
 form of the results, that the particles are identical
 and of the same mass. In addition to this we
 must assume that the particles are in a state of
 equilibrium. The results of the theory are
 therefore for a system of three or more particles
 which are in a state of equilibrium, and the results
 are identical with those for a system of three
 particles. The results of the theory are
 therefore for a system of three or more particles
 which are in a state of equilibrium, and the results
 are identical with those for a system of three
 particles.

$$m_I = \frac{e}{2Mc} \cdot I \cdot \frac{e}{2Mc}$$

The results of the theory are therefore for a system of three or more particles which are in a state of equilibrium, and the results are identical with those for a system of three particles.

$$\left(\frac{e}{2Mc}\right) I = \left(\frac{e}{2Mc}\right) I$$

The results of the theory are therefore for a system of three or more particles which are in a state of equilibrium, and the results are identical with those for a system of three particles.

$$\left(\frac{e}{2Mc}\right) M = \left(\frac{e}{2Mc}\right) M$$

The results of the theory are therefore for a system of three or more particles which are in a state of equilibrium, and the results are identical with those for a system of three particles.

It is true that the results of the theory are for a system of three or more particles which are in a state of equilibrium, and the results are identical with those for a system of three particles.

there is a torque \underline{L} exerted on the dipole given by the expression¹⁷

$$\underline{L} = \mu \times \underline{H}.$$

17. Pake, G.E. Am. Jour. Phys. 18, 438 (1950).

Since the rate of change of angular momentum, \underline{p} , of a system is equal to the applied torque, there results

$$\frac{d\underline{p}}{dt} = \underline{L},$$

or

$$\frac{d\underline{p}}{dt} = \mu \times \underline{H}.$$

The vector expression for the nuclear magnetic moment is

$$\mu_I = g_I \cdot \frac{e}{2mc} \cdot \underline{p}_I,$$

so for a nuclear dipole in a magnetic field one may write

$$\frac{d\underline{p}_I}{dt} = -g_I \cdot \frac{e}{2mc} \cdot \underline{H} \times \underline{p}_I.$$

From classical mechanics we also have the vector relationship

$$\frac{d\underline{p}}{dt} = \underline{\omega} \times \underline{p},$$

so it may be concluded that if a nucleus of magnetic moment vector $\mu_I = g_I \cdot \frac{e}{2mc} \cdot \underline{p}_I$ is placed in a magnetic field, the magnetic moment vector of the nucleus will precess about the direction of \underline{H} with the angular frequency

$$\omega_0 = -g_I \cdot \frac{e}{2mc} \cdot \underline{H}.$$

This frequency of precession is called the Larmor precession

frequency. From the last equation it is noted that ω_0 is not dependent on the angle between μ_I and \underline{H} .

Therefore, the total energy is the sum of the kinetic and potential energies.

$$E = K + U$$

12. The total energy is the sum of the kinetic and potential energies.

13. The total energy is the sum of the kinetic and potential energies.

14. The total energy is the sum of the kinetic and potential energies.

$$E = K + U$$

$$E = K + U$$

15. The total energy is the sum of the kinetic and potential energies.

$$E = K + U$$

16. The total energy is the sum of the kinetic and potential energies.

$$E = K + U$$

17. The total energy is the sum of the kinetic and potential energies.

$$E = K + U$$

18. The total energy is the sum of the kinetic and potential energies.

19. The total energy is the sum of the kinetic and potential energies.

20. The total energy is the sum of the kinetic and potential energies.

21. The total energy is the sum of the kinetic and potential energies.

$$E = K + U$$

22. The total energy is the sum of the kinetic and potential energies.

23. The total energy is the sum of the kinetic and potential energies.

$$E = K + U$$

The potential energy U of a magnetic dipole μ in a field H is $U = |\mu||H|(1 - \cos \theta)$, where θ is the angle between μ and H . This expression gives a maximum for U of $2|\mu||H|$ when μ and H are antiparallel. Since the nuclear angular momentum vector may assume $2I+1$ orientations with respect to H , with $M_I = I, I-1, \dots, -I+1, -I$, there will be in all $2I$ energy levels due to nuclear spin orientations, each differing in energy by

$$\frac{2(\mu_H)_{\max} H}{2I} \quad \text{or} \quad \frac{(\mu_H)_{\max} H}{I} \text{ ergs.}$$

The selection rule for transitions between the $2I+1$ possible orientations of the magnetic moment vector is

$$\Delta M_I = \pm 1.$$

This requires that the nucleus emit or absorb

$$\Delta E = h\nu = \frac{(\mu_H)_{\max} H}{I} \text{ ergs.}$$

in event of change of orientation of spin vector. The frequency obtained from this expression is seen to be

$$\nu = \frac{(\mu_H)_{\max} H}{Ih}.$$

But

$$\frac{(\mu_H)_{\max} H}{Ih} = g_I \cdot \frac{e}{2Mc} \cdot I \frac{h}{2\pi} \cdot \frac{1}{Ih},$$

or

$$\nu = g_I \cdot \frac{e}{4\pi Mc} \cdot H$$

Using $\omega = 2\pi\nu$ one obtains

$$\omega = g_I \cdot \frac{e}{2Mc} \cdot H,$$

which is the absolute value previously found for the Larmor angular precession frequency. In other words, an oscillating electromagnetic field of frequency

$$\nu = g_I \cdot \frac{e}{4\pi Mc} \cdot H$$

The potential energy U of a dipole in an electric field E is given by

$$U = -p \cdot E = -pE \cos \theta \quad (1)$$

This expression gives a relation for U of p and E when θ is known.

For equilibrium, the net torque on the dipole must be zero.

Let the dipole be in equilibrium at an angle θ with the electric field E .

The forces acting on the dipole are the electric forces qE and $-qE$.

The torque due to these forces is given by

$$\tau = \frac{qE \cdot 2a \sin \theta}{2} = qEa \sin \theta$$

The electric field E is related to the potential U by

$E = -\frac{dU}{dr}$

$$E = \frac{U}{r}$$

This relation can be used to find the torque on the dipole.

$$\tau = \frac{qE \cdot 2a \sin \theta}{2} = qEa \sin \theta$$

At equilibrium, the net torque on the dipole is zero.

Thus, the dipole is in equilibrium when $\theta = 0$ or $\theta = \pi$.

$$\frac{qE \cdot 2a \sin \theta}{2} = 0$$

$$\frac{1}{2} \cdot \frac{qE \cdot 2a \sin \theta}{2} = \frac{qE \cdot 2a \sin \theta}{2}$$

$$\frac{1}{2} \cdot \frac{qE \cdot 2a \sin \theta}{2} = \frac{qE \cdot 2a \sin \theta}{2}$$

where $\omega = 2\pi \nu$ is the angular frequency.

$$\omega = \frac{qE \cdot 2a \sin \theta}{2}$$

where ω is the angular frequency of the dipole's oscillation.

The electric field E is related to the potential U by

$$E = -\frac{dU}{dr}$$

applied to a nucleus in the external magnetic field H might be capable of inducing transitions between adjacent nuclear spin orientation states. In the following four sections there will be discussed the four most recent methods developed for applying the above principle.

All of the four methods depend upon the Maxwell-Boltzmann relation that, in any system in thermal equilibrium, the ratio of the populations of states with different energies, say E_1 and E_2 , is

$$\frac{N(E_1)}{N(E_2)} = \frac{e^{-E_1/kT}}{e^{-E_2/kT}}$$

If this relation is applied to the case of protons in a magnetic field H , the ratio of the populations of the two proton states $M_I = +\frac{1}{2}$ and $M_I = -\frac{1}{2}$ is seen to be

$$\frac{N(+\frac{1}{2})}{N(-\frac{1}{2})} = \frac{e^{-E_1/kT}}{e^{-E_2/kT}} = e^{(E_2-E_1)/kT} = e^{\Delta E/kT},$$

in which $(E_2 - E_1) = \Delta E = g_I \mu_n H$.

For temperature $T = 300^\circ\text{K}$ and magnetic field strengths available in laboratories $\frac{\Delta E}{kT}$ is small, so the last equation may be written

$$\frac{N(+\frac{1}{2})}{N(-\frac{1}{2})} \cong 1 + g_I \mu_n \frac{H}{kT}.$$

For a field of 10,000 gauss, $g_I = 5.58$ for protons, and $kT = 4 \times 10^{-14}$ erg the last expression gives

$$\frac{N(+\frac{1}{2})}{N(-\frac{1}{2})} \cong 1 + 7 \times 10^{-6}.$$

This indicates that for each million protons in the higher energy state there are one million and seven protons in the state

$$\frac{N(E_1)}{N(E_2)} = \frac{e^{-\frac{E_1}{kT}}}{e^{-\frac{E_2}{kT}}}$$

$$\frac{N(E_1)}{N(E_2)} = \frac{e^{-\frac{E_1}{kT}}}{e^{-\frac{E_2}{kT}}} = \frac{N(E_1)}{N(E_2)}$$

$$H_{\text{max}} = -k \ln \left(\frac{1}{2} \right) = k \ln 2$$

$$\frac{N(E_1)}{N(E_2)} = \frac{e^{-\frac{E_1}{kT}}}{e^{-\frac{E_2}{kT}}}$$

$$\frac{N(E_1)}{N(E_2)} = \frac{e^{-\frac{E_1}{kT}}}{e^{-\frac{E_2}{kT}}}$$

with lower energy. Clearly this relationship may be extended to nuclei with spin I possibly other than $\frac{1}{2}$. The results of the above example will be referred to in some of the forthcoming discussions.

Another important factor in the methods to be discussed is the spin-lattice relaxation time T_1 for the nucleus of interest in a given liquid or solid. Thermodynamically, a relaxation process in the case of nuclei is any method of energy exchange between the system of nuclear spins and the lattice. Thus if a sample is placed suddenly into a magnetic field it is important for the experimenter to know how long a period of time on the average will elapse before the equilibrium number of nuclei will be found in the upper and lower energy states. T_1 is employed as a measure of this time and is defined as the length of time required for all except the fraction $(1/e)$ of the equilibrium excess number of nuclei to reach the lower energy state. Relaxation mechanisms in fluids have been investigated by Purcell and his group¹⁸ at Harvard who have determined that

18. Bloembergen, N., Purcell, E.M., and Pound, R.V. Phys. Rev. **73**, 679 (1948).

Brownian motions at the Larmor frequency are responsible for the processes. In liquids T_1 ordinarily decreases with increase of viscosity. In some cases, however, T_1 decreases with increasing viscosity until a minimum value is reached and after that it increases with increase in viscosity.

with some energy. Clearly this relationship may be extended to
 model with this. I possibly order than 1. The results of the above
 analysis will be reported in some of the following discussion.
 Another important factor in the system to be discussed in the
 only-factor relationship than 1. For the system of interest in a
 given field or field. (Theoretically, a relationship between the
 the case of model is the method of using evidence between the sys-
 tem of analysis system and the system. That is a model is a model
 whereby the a system field is a model field. The relationship
 to form the field a model of time in the system will be the model
 the relationship model of model will be model to the system and lower
 every model. It is model as a model of this field and is
 defined as the model of the model for all other the model
 (1/2) of the relationship model model of model to model the model
 every model. Relationship model in this field model (model)
 to model and the model a model the model between model

The relationship of model, 1.1 and model, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7.0, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9, 8.0, 8.1, 8.2, 8.3, 8.4, 8.5, 8.6, 8.7, 8.8, 8.9, 9.0, 9.1, 9.2, 9.3, 9.4, 9.5, 9.6, 9.7, 9.8, 9.9, 10.0, 10.1, 10.2, 10.3, 10.4, 10.5, 10.6, 10.7, 10.8, 10.9, 11.0, 11.1, 11.2, 11.3, 11.4, 11.5, 11.6, 11.7, 11.8, 11.9, 12.0, 12.1, 12.2, 12.3, 12.4, 12.5, 12.6, 12.7, 12.8, 12.9, 13.0, 13.1, 13.2, 13.3, 13.4, 13.5, 13.6, 13.7, 13.8, 13.9, 14.0, 14.1, 14.2, 14.3, 14.4, 14.5, 14.6, 14.7, 14.8, 14.9, 15.0, 15.1, 15.2, 15.3, 15.4, 15.5, 15.6, 15.7, 15.8, 15.9, 16.0, 16.1, 16.2, 16.3, 16.4, 16.5, 16.6, 16.7, 16.8, 16.9, 17.0, 17.1, 17.2, 17.3, 17.4, 17.5, 17.6, 17.7, 17.8, 17.9, 18.0, 18.1, 18.2, 18.3, 18.4, 18.5, 18.6, 18.7, 18.8, 18.9, 19.0, 19.1, 19.2, 19.3, 19.4, 19.5, 19.6, 19.7, 19.8, 19.9, 20.0, 20.1, 20.2, 20.3, 20.4, 20.5, 20.6, 20.7, 20.8, 20.9, 21.0, 21.1, 21.2, 21.3, 21.4, 21.5, 21.6, 21.7, 21.8, 21.9, 22.0, 22.1, 22.2, 22.3, 22.4, 22.5, 22.6, 22.7, 22.8, 22.9, 23.0, 23.1, 23.2, 23.3, 23.4, 23.5, 23.6, 23.7, 23.8, 23.9, 24.0, 24.1, 24.2, 24.3, 24.4, 24.5, 24.6, 24.7, 24.8, 24.9, 25.0, 25.1, 25.2, 25.3, 25.4, 25.5, 25.6, 25.7, 25.8, 25.9, 26.0, 26.1, 26.2, 26.3, 26.4, 26.5, 26.6, 26.7, 26.8, 26.9, 27.0, 27.1, 27.2, 27.3, 27.4, 27.5, 27.6, 27.7, 27.8, 27.9, 28.0, 28.1, 28.2, 28.3, 28.4, 28.5, 28.6, 28.7, 28.8, 28.9, 29.0, 29.1, 29.2, 29.3, 29.4, 29.5, 29.6, 29.7, 29.8, 29.9, 30.0, 30.1, 30.2, 30.3, 30.4, 30.5, 30.6, 30.7, 30.8, 30.9, 31.0, 31.1, 31.2, 31.3, 31.4, 31.5, 31.6, 31.7, 31.8, 31.9, 32.0, 32.1, 32.2, 32.3, 32.4, 32.5, 32.6, 32.7, 32.8, 32.9, 33.0, 33.1, 33.2, 33.3, 33.4, 33.5, 33.6, 33.7, 33.8, 33.9, 34.0, 34.1, 34.2, 34.3, 34.4, 34.5, 34.6, 34.7, 34.8, 34.9, 35.0, 35.1, 35.2, 35.3, 35.4, 35.5, 35.6, 35.7, 35.8, 35.9, 36.0, 36.1, 36.2, 36.3, 36.4, 36.5, 36.6, 36.7, 36.8, 36.9, 37.0, 37.1, 37.2, 37.3, 37.4, 37.5, 37.6, 37.7, 37.8, 37.9, 38.0, 38.1, 38.2, 38.3, 38.4, 38.5, 38.6, 38.7, 38.8, 38.9, 39.0, 39.1, 39.2, 39.3, 39.4, 39.5, 39.6, 39.7, 39.8, 39.9, 40.0, 40.1, 40.2, 40.3, 40.4, 40.5, 40.6, 40.7, 40.8, 40.9, 41.0, 41.1, 41.2, 41.3, 41.4, 41.5, 41.6, 41.7, 41.8, 41.9, 42.0, 42.1, 42.2, 42.3, 42.4, 42.5, 42.6, 42.7, 42.8, 42.9, 43.0, 43.1, 43.2, 43.3, 43.4, 43.5, 43.6, 43.7, 43.8, 43.9, 44.0, 44.1, 44.2, 44.3, 44.4, 44.5, 44.6, 44.7, 44.8, 44.9, 45.0, 45.1, 45.2, 45.3, 45.4, 45.5, 45.6, 45.7, 45.8, 45.9, 46.0, 46.1, 46.2, 46.3, 46.4, 46.5, 46.6, 46.7, 46.8, 46.9, 47.0, 47.1, 47.2, 47.3, 47.4, 47.5, 47.6, 47.7, 47.8, 47.9, 48.0, 48.1, 48.2, 48.3, 48.4, 48.5, 48.6, 48.7, 48.8, 48.9, 49.0, 49.1, 49.2, 49.3, 49.4, 49.5, 49.6, 49.7, 49.8, 49.9, 50.0, 50.1, 50.2, 50.3, 50.4, 50.5, 50.6, 50.7, 50.8, 50.9, 51.0, 51.1, 51.2, 51.3, 51.4, 51.5, 51.6, 51.7, 51.8, 51.9, 52.0, 52.1, 52.2, 52.3, 52.4, 52.5, 52.6, 52.7, 52.8, 52.9, 53.0, 53.1, 53.2, 53.3, 53.4, 53.5, 53.6, 53.7, 53.8, 53.9, 54.0, 54.1, 54.2, 54.3, 54.4, 54.5, 54.6, 54.7, 54.8, 54.9, 55.0, 55.1, 55.2, 55.3, 55.4, 55.5, 55.6, 55.7, 55.8, 55.9, 56.0, 56.1, 56.2, 56.3, 56.4, 56.5, 56.6, 56.7, 56.8, 56.9, 57.0, 57.1, 57.2, 57.3, 57.4, 57.5, 57.6, 57.7, 57.8, 57.9, 58.0, 58.1, 58.2, 58.3, 58.4, 58.5, 58.6, 58.7, 58.8, 58.9, 59.0, 59.1, 59.2, 59.3, 59.4, 59.5, 59.6, 59.7, 59.8, 59.9, 60.0, 60.1, 60.2, 60.3, 60.4, 60.5, 60.6, 60.7, 60.8, 60.9, 61.0, 61.1, 61.2, 61.3, 61.4, 61.5, 61.6, 61.7, 61.8, 61.9, 62.0, 62.1, 62.2, 62.3, 62.4, 62.5, 62.6, 62.7, 62.8, 62.9, 63.0, 63.1, 63.2, 63.3, 63.4, 63.5, 63.6, 63.7, 63.8, 63.9, 64.0, 64.1, 64.2, 64.3, 64.4, 64.5, 64.6, 64.7, 64.8, 64.9, 65.0, 65.1, 65.2, 65.3, 65.4, 65.5, 65.6, 65.7, 65.8, 65.9, 66.0, 66.1, 66.2, 66.3, 66.4, 66.5, 66.6, 66.7, 66.8, 66.9, 67.0, 67.1, 67.2, 67.3, 67.4, 67.5, 67.6, 67.7, 67.8, 67.9, 68.0, 68.1, 68.2, 68.3, 68.4, 68.5, 68.6, 68.7, 68.8, 68.9, 69.0, 69.1, 69.2, 69.3, 69.4, 69.5, 69.6, 69.7, 69.8, 69.9, 70.0, 70.1, 70.2, 70.3, 70.4, 70.5, 70.6, 70.7, 70.8, 70.9, 71.0, 71.1, 71.2, 71.3, 71.4, 71.5, 71.6, 71.7, 71.8, 71.9, 72.0, 72.1, 72.2, 72.3, 72.4, 72.5, 72.6, 72.7, 72.8, 72.9, 73.0, 73.1, 73.2, 73.3, 73.4, 73.5, 73.6, 73.7, 73.8, 73.9, 74.0, 74.1, 74.2, 74.3, 74.4, 74.5, 74.6, 74.7, 74.8, 74.9, 75.0, 75.1, 75.2, 75.3, 75.4, 75.5, 75.6, 75.7, 75.8, 75.9, 76.0, 76.1, 76.2, 76.3, 76.4, 76.5, 76.6, 76.7, 76.8, 76.9, 77.0, 77.1, 77.2, 77.3, 77.4, 77.5, 77.6, 77.7, 77.8, 77.9, 78.0, 78.1, 78.2, 78.3, 78.4, 78.5, 78.6, 78.7, 78.8, 78.9, 79.0, 79.1, 79.2, 79.3, 79.4, 79.5, 79.6, 79.7, 79.8, 79.9, 80.0, 80.1, 80.2, 80.3, 80.4, 80.5, 80.6, 80.7, 80.8, 80.9, 81.0, 81.1, 81.2, 81.3, 81.4, 81.5, 81.6, 81.7, 81.8, 81.9, 82.0, 82.1, 82.2, 82.3, 82.4, 82.5, 82.6, 82.7, 82.8, 82.9, 83.0, 83.1, 83.2, 83.3, 83.4, 83.5, 83.6, 83.7, 83.8, 83.9, 84.0, 84.1, 84.2, 84.3, 84.4, 84.5, 84.6, 84.7, 84.8, 84.9, 85.0, 85.1, 85.2, 85.3, 85.4, 85.5, 85.6, 85.7, 85.8, 85.9, 86.0, 86.1, 86.2, 86.3, 86.4, 86.5, 86.6, 86.7, 86.8, 86.9, 87.0, 87.1, 87.2, 87.3, 87.4, 87.5, 87.6, 87.7, 87.8, 87.9, 88.0, 88.1, 88.2, 88.3, 88.4, 88.5, 88.6, 88.7, 88.8, 88.9, 89.0, 89.1, 89.2, 89.3, 89.4, 89.5, 89.6, 89.7, 89.8, 89.9, 90.0, 90.1, 90.2, 90.3, 90.4, 90.5, 90.6, 90.7, 90.8, 90.9, 91.0, 91.1, 91.2, 91.3, 91.4, 91.5, 91.6, 91.7, 91.8, 91.9, 92.0, 92.1, 92.2, 92.3, 92.4, 92.5, 92.6, 92.7, 92.8, 92.9, 93.0, 93.1, 93.2, 93.3, 93.4, 93.5, 93.6, 93.7, 93.8, 93.9, 94.0, 94.1, 94.2, 94.3, 94.4, 94.5, 94.6, 94.7, 94.8, 94.9, 95.0, 95.1, 95.2, 95.3, 95.4, 95.5, 95.6, 95.7, 95.8, 95.9, 96.0, 96.1, 96.2, 96.3, 96.4, 96.5, 96.6, 96.7, 96.8, 96.9, 97.0, 97.1, 97.2, 97.3, 97.4, 97.5, 97.6, 97.7, 97.8, 97.9, 98.0, 98.1, 98.2, 98.3, 98.4, 98.5, 98.6, 98.7, 98.8, 98.9, 99.0, 99.1, 99.2, 99.3, 99.4, 99.5, 99.6, 99.7, 99.8, 99.9, 100.0, 100.1, 100.2, 100.3, 100.4, 100.5, 100.6, 100.7, 100.8, 100.9, 101.0, 101.1, 101.2, 101.3, 101.4, 101.5, 101.6, 101.7, 101.8, 101.9, 102.0, 102.1, 102.2, 102.3, 102.4, 102.5, 102.6, 102.7, 102.8, 102.9, 103.0, 103.1, 103.2, 103.3, 103.4, 103.5, 103.6, 103.7, 103.8, 103.9, 104.0, 104.1, 104.2, 104.3, 104.4, 104.5, 104.6, 104.7, 104.8, 104.9, 105.0, 105.1, 105.2, 105.3, 105.4, 105.5, 105.6, 105.7, 105.8, 105.9, 106.0, 106.1, 106.2, 106.3, 106.4, 106.5, 106.6, 106.7, 106.8, 106.9, 107.0, 107.1, 107.2, 107.3, 107.4, 107.5, 107.6, 107.7, 107.8, 107.9, 108.0, 108.1, 108.2, 108.3, 108.4, 108.5, 108.6, 108.7, 108.8, 108.9, 109.0, 109.1, 109.2, 109.3, 109.4, 109.5, 109.6, 109.7, 109.8, 109.9, 110.0, 110.1, 110.2, 110.3, 110.4, 110.5, 110.6, 110.7, 110.8, 110.9, 111.0, 111.1, 111.2, 111.3, 111.4, 111.5, 111.6, 111.7, 111.8, 111.9, 112.0, 112.1, 112.2, 112.3, 112.4, 112.5, 112.6, 112.7, 112.8, 112.9, 113.0, 113.1, 113.2, 113.3, 113.4, 113.5, 113.6, 113.7, 113.8, 113.9, 114.0, 114.1, 114.2, 114.3, 114.4, 114.5, 114.6, 114.7, 114.8, 114.9, 115.0, 115.1, 115.2, 115.3, 115.4, 115.5, 115.6, 115.7, 115.8, 115.9, 116.0, 116.1, 116.2, 116.3, 116.4, 116.5, 116.6, 116.7, 116.8, 116.9, 117.0, 117.1, 117.2, 117.3, 117.4, 117.5, 117.6, 117.7, 117.8, 117.9, 118.0, 118.1, 118.2, 118.3, 118.4, 118.5, 118.6, 118.7, 118.8, 118.9, 119.0, 119.1, 119.2, 119.3, 119.4, 119.5, 119.6, 119.7, 119.8, 119.9, 120.0, 120.1, 120.2, 120.3, 120.4, 120.5, 120.6, 120.7, 120.8, 120.9, 121.0, 121.1, 121.2, 121.3, 121.4, 121.5, 121.6, 121.7, 121.8, 121.9, 122.0, 122.1, 122.2, 122.3, 122.4, 122.5, 122.6, 122.7, 122.8, 122.9, 123.0, 123.1, 123.2, 123.3, 123.4, 123.5, 123.6, 123.7, 123.8, 123.9, 124.0, 124.1, 124.2, 124.3, 124.4, 124.5, 124.6, 124.7, 124.8, 124.9, 125.0, 125.1, 125.2, 125.3, 125.4, 125.5, 125.6, 125.7, 125.8, 125.9, 126.0, 126.1, 126.2, 126.3, 126.4, 126.5, 126.6, 126.7, 126.8, 126.9, 127.0, 127.1, 127.2, 127.3, 127.4, 127.5, 127.6, 127.7, 127.8, 127.9, 128.0, 128.1, 128.2, 128.3, 128.4, 128.5, 128.6, 128.7, 128.8, 128.9, 129.0, 129.1, 129.2, 129.3, 129.4, 129.5, 129.6, 129.7, 129.8, 129.9, 130.0, 130.1, 130.2, 130.3, 130.4, 130.5, 130.6, 130.7, 130.8, 130.9, 131.0, 131.1, 131.2, 131.3, 131.4, 131.5, 131.6, 131.7, 131.8, 131.9, 132.0, 132.1, 132.2, 132.3, 132.4, 132.5, 132.6, 132.7, 132.8, 132.9, 133.0, 133.1, 133.2, 133.3, 133.4, 133.5, 133.6, 133.7, 133.8, 133.9, 134.0, 134.1, 134.2, 134.3, 134.4, 134.5, 134.6, 134.7, 134.8, 134.9, 135.0, 135.1, 135.2, 135.3, 135.4, 135.5, 135.6, 135.7, 135.8, 135.9, 136.0, 136.1, 136.2, 136.3, 136.4, 136.5, 136.6, 136.7, 136.8, 136.9, 137.0, 137.1, 137.2, 137.3, 137.4, 137.5, 137.6, 137.7, 137.8, 137.9, 138.0, 138.1, 138.2, 138.3, 138.4, 138.5, 138.6, 138.7, 138.8, 138.9, 139.0, 139.1, 139.2, 139.3, 139.4, 139.5, 139.6, 139.7, 139.8, 139.9, 140.0, 140.1, 140.2, 140.3, 140.4, 140.5, 140.6, 140.7, 140.8, 140.9, 141.0, 141.1, 141.2, 141.3, 141.4, 141.5, 141.6, 141.7, 141.8, 141.9, 142.0, 142.1, 142.2, 142.3, 142.4, 142.5, 142.6, 142.7, 142.8, 142.9, 143.0, 143.1, 143.2, 143.3, 143.4, 143.5, 143.6, 143.7, 143.8, 143.9, 144.0, 144.1, 144.2, 144.3, 144.4, 144.5, 144.6, 144.7, 144.8, 144.9, 145.0, 145.1, 145.2, 145.3, 145.4, 145.5, 145.6, 145.7, 145.8, 145.9, 146.0, 146.1, 146.2, 146.3, 146.4, 146.5, 146.6, 146.7, 146.8, 146.9, 147.0, 147.1, 147.2, 147.3, 147.4, 147.5, 147.6, 147.7, 147.8, 147.9, 148.0, 148.1, 148.2, 148.3, 148.4, 148.5, 148.6, 148.7, 148.8, 148.9, 149.0, 149.1, 149.2, 149.3, 149.4, 149.5, 149.6, 149.7, 149.8, 149.9, 150.0, 150.1, 150.2, 150.3, 150.4, 150.5, 150.6, 150.7, 150.8, 150.9, 151.0, 151.1, 151.2, 151.3, 151.4, 151.5, 151.6, 151.7, 151.8, 151.9, 152.0, 152.1, 152.2, 152.3, 152.4, 152.5, 152.6, 152.7, 152.8, 152.9, 153.0, 153.1, 153.2, 153.3, 153.4, 153.5, 153.6, 153.7, 153.8, 153.9, 154.0, 154.1, 154.2, 154.3, 154.4, 154.5, 154.6, 154.7, 154.8, 154.9, 155.0, 155.1, 155.2, 155.3, 155.4, 155.5, 155.6, 155.7, 155.8, 155.9, 156.0, 156.1, 156.2, 156.3, 156.4, 156.5, 156.6, 156.7, 156.8, 156.9, 157.0, 157.1, 157.2, 157.3, 157.4, 157.5, 157.6, 157.7, 157.8, 157.9, 158.0, 158.1, 158.2, 158.3, 158.4, 158.5, 158.6, 158.7, 158.8, 158.9, 159.0, 159.1, 159.2, 159.3, 159.4, 159.5, 159.6, 159.7, 159.8, 159.9, 160.0, 160.1, 160.2, 160.3, 160.4, 160.5, 160.6, 160.7, 160.8, 160.9, 161.0, 161.1, 161.2, 161.3, 161.4, 161.5, 161.6, 161.7, 161.8, 161.9, 162.0, 162.1, 162.2, 162.3, 162.4, 162.5, 162.6, 162.7, 162.8, 162.9, 163.0, 163.1, 163.2, 163.3, 163.4, 163.5, 163.6, 163.7, 163.8, 163.9, 164.0, 164.1, 164.2, 164.3, 164.4, 164.5, 164.6, 164.7, 164.8, 164.9, 165.0, 165.1, 165.2, 165.3, 165.4, 165.5, 165.6, 165.7, 165.8, 165.9, 166.0, 166.1, 166.2, 166.3, 166.4, 166.5, 166.6, 166.7, 166.8, 166.9, 167.0, 167.1, 167.2, 167.3, 167.4, 167.5, 167.6, 167.7, 167.8, 167.9, 168.0, 168.1, 168.2, 168.3, 168.4, 168.5, 168.6, 168.7, 168.8, 168.9, 169.0, 169.1, 169.2, 169.3, 169.4, 169.5, 169.6, 169.7, 169.8, 169.9, 170.0, 170.1, 170.2, 170.3, 170.4, 170.5, 170.6, 170.7, 170.8, 170.9, 171.0, 171.1, 171.2, 171.3, 171.4, 171.5, 171.6, 171.7, 171.8, 171.9, 172.0, 172.1, 172.2, 172.3, 172.4, 172.5, 172.6, 172.7, 172.8, 172.9, 173.0, 173.1, 173.2, 173.3, 173.4, 173.5, 173.6, 173.7, 173.8, 173.9, 174.0, 174.1, 174.2, 174.3, 174.4, 174.5, 174.6, 174.7, 174.8, 174.9, 175.0, 175.1, 175.2, 175.3, 175.4, 175.5, 175.6, 175.7, 175.8, 175.9, 176.0, 176.1, 176.2, 176.3, 176.4, 176.5, 176.6, 176.7, 176.8, 176.9, 177.0, 177.1, 177.2, 177.3, 177.4, 177.5, 177.6, 177.7, 177.8, 177.9, 178.0, 178.1, 178.2, 178.3, 178.4, 178.5, 178.6, 178.7, 178.8, 178.9, 179.0, 179.1, 179.2, 179.3, 179.4, 179.5, 179.6, 179.7, 179.8, 179.9, 180.0, 180.1, 180.2, 180.3, 180.4, 180.5, 180.6, 180.7, 180.8, 180.9, 181.0, 181.1, 181.2, 181.3, 181.4, 181.5, 181.6, 181.7, 181.8, 181.9, 182.0, 182.1, 182.2, 182.3, 182.4, 182.5, 182.6, 182.7, 182.8, 182.9, 183.0, 183.1, 183.2, 183.3, 183.4, 183.5, 183.6, 183.7, 183.8, 183.9, 184.0, 184.1, 184.2, 184.3, 184.4, 184.5, 184.6, 184.7, 184.8, 184.9, 185.0, 185.1, 185.2, 185.3, 185.4, 185.5, 185.6, 185.7, 185.8, 185.9, 186.0, 186.1, 186.2, 186.3, 186.4, 186.5, 186.6, 186.7, 186.8, 186.9, 187.0, 187.1, 187.2, 187.3, 187.4, 187.5, 187.6, 187.7, 187.8, 187.9, 188.0, 188.1, 188.2, 188.3

The addition of paramagnetic ions to a sample frequently will give a marked decrease in T_1 by increasing the local perturbing effects upon nuclei of the sample.

(1) Method of Purcell and Pound

In the method developed by Purcell, Pound, and Torrey,¹⁸ nuclear absorption unbalances an r-f bridge, giving rise to a signal in a receiver used as a detector. The sensitivity of this method is very good, being as high perhaps as any of the other three methods discussed in this section. Purcell has demonstrated that nuclear resonance with this method can be observed in a sample containing only 10^{19} atoms.

In this method a signal generator is employed as the r-f signal source for the bridge. An essential feature of the bridge is the reduction effected in the relative magnitude of output fluctuations arising from amplitude fluctuations in the signal supplied by the r-f signal generator. This is done by obtaining a voltage node in the bridge output to the amplifier when the bridge is placed in balance with the system not adjusted for nuclear resonance. One method to effect this voltage node has been to place an extra half-wave-length of cable in one arm of the bridge. Another important feature of this method is the reduction in the r-f level at the input to the amplifier to permit considerable r-f amplification before detection.

One arm of the bridge contains a tuned circuit having in the gap between the pole faces of an electric magnet a coil into which a

[illegible][illegible]

THE NATIONAL ARCHIVES HAS A STRONG INTEREST IN THE PRESERVATION OF THE RECORDS OF THE PRESIDENTS OF THE UNITED STATES.

sample to be studied can be inserted. In the other arm of the bridge is a dummy circuit similar in all respects to the other but not having its coil located in the magnetic field. Nuclear resonance absorption in the sample causes a change in the balance of the r-f bridge which is detected by means of the receiver. When liquids of high dielectric constants are being studied a dummy sample is sometimes placed in the dummy coil to balance as nearly as possible the appreciable change in stray capacitance resulting from the sample being placed in the coil of the other arm.

The strong magnetic field H_0 is modulated sinusoidally at 30 cycles per second by auxiliary coils mounted on the pole pieces. This modulation is parallel to the main field with peak strength never more than 15 gauss. Near resonance the low frequency modulation of the magnetic field causes the Larmor frequency of nuclear precession to vary in and out of resonance with the result that resonance absorption occurs in the arm of the bridge containing the sample and the bridge is thrown out of balance. The signal so produced is amplified and suitably detected.

An improvement in the above technique has been the elimination of the half-wave-length cable, which is several meters long and is frequently the cause for considerable troublesome instability.

With properly adjusted equipment operating under suitable conditions the sensitivity of this method depends on the width of the observed resonance. The natural width of the resonance signal depends on the substance used and in some liquids is less than a thousandth of a gauss. This may cause the homogeneity of the magnetic field to be a limiting factor in the sensitivity.

...the ... of the ...

The r-f bridge nuclear resonance spectrometer is well suited for the observation of shapes and widths of absorption lines in various substances. Four such resonance absorption lines for protons in different substances which were obtained by Purcell using the bridge method are shown in Figure 2.¹⁹ The line displayed in Figure 2(a)

19. Purcell, E.M. Science 107, 433 (1948). ABSORPTION LINE

was obtained from protons in water. This absorption occurred within an interval of only 0.2 gauss in a field of 6,940 gauss. Other substances containing protons display an absorption line at the same field strength if the radio frequency is the same, but the line width varies greatly with different substances. The proton resonance in ice, shown in Figure 2(b), is actually 20 to 50 times as wide as the line obtained in water. In other crystals the variation in line width is even more striking and a pronounced difference in line shapes is observed as one may notice by comparing the various absorption lines shown in this figure.

The variations in line widths and line shapes find no explanation from nuclear properties but are associated with the perturbing magnetic fields which have their origins in the immediate surroundings of a given hydrogen atom and depend on both the location and motion of its immediate neighbors. It perhaps seems strange, but the sharpest resonance lines are obtained from liquids, whose molecules compared to those of solids are in a greater state of random motion. This random thermal motion of the molecules occurs at a

THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION

THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION
455 N. 5TH ST. NEW YORK, N. Y. 10017
ASTOR LENOX TILDEN FOUNDATION
15 EAST 17TH ST. NEW YORK, N. Y. 10003

THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION

THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION
455 N. 5TH ST. NEW YORK, N. Y. 10017
ASTOR LENOX TILDEN FOUNDATION
15 EAST 17TH ST. NEW YORK, N. Y. 10003
THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION
455 N. 5TH ST. NEW YORK, N. Y. 10017
ASTOR LENOX TILDEN FOUNDATION
15 EAST 17TH ST. NEW YORK, N. Y. 10003
THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION
455 N. 5TH ST. NEW YORK, N. Y. 10017
ASTOR LENOX TILDEN FOUNDATION
15 EAST 17TH ST. NEW YORK, N. Y. 10003
THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION
455 N. 5TH ST. NEW YORK, N. Y. 10017
ASTOR LENOX TILDEN FOUNDATION
15 EAST 17TH ST. NEW YORK, N. Y. 10003

THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION
455 N. 5TH ST. NEW YORK, N. Y. 10017
ASTOR LENOX TILDEN FOUNDATION
15 EAST 17TH ST. NEW YORK, N. Y. 10003
THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION
455 N. 5TH ST. NEW YORK, N. Y. 10017
ASTOR LENOX TILDEN FOUNDATION
15 EAST 17TH ST. NEW YORK, N. Y. 10003
THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION
455 N. 5TH ST. NEW YORK, N. Y. 10017
ASTOR LENOX TILDEN FOUNDATION
15 EAST 17TH ST. NEW YORK, N. Y. 10003
THE NEW YORK PUBLIC LIBRARY ASTOR LENOX TILDEN FOUNDATION
455 N. 5TH ST. NEW YORK, N. Y. 10017
ASTOR LENOX TILDEN FOUNDATION
15 EAST 17TH ST. NEW YORK, N. Y. 10003

FIGURE 2 (a).

PROTON RESONANCE
ABSORPTION CURVE




FIGURE 2 (b).

(b). IN ICE.



FIGURE 2 (c).

(c). IN POWDERED
GYPSUM.





FIGURE 2 (d).

(d). IN SINGLE
GYPSUM
CRYSTAL.



Curves Not
to Same
Scale.
(After Purcell '9)

frequency higher than that of the proton's nuclear precession. It has been shown statistically that the very rapidity of this thermal motion greatly reduces the perturbing effect from neighboring atoms, so that local perturbing effects on the applied magnetic field are much less than in the case of solid substances.¹³

In the case of a single crystal of gypsum referred to in Figure 2(d), protons in different parts of the molecule have different, more nearly constant local perturbing fields which roughly are the same on corresponding protons in different molecules. This results in a broad line with four resonance peaks corresponding to four different values of field strength observed.

In the case of powdered gypsum a greater degree of randomness of orientation of molecules within the sample occurs. The statistical effect of this (imperfect) randomness is to cause the absorption curve to exhibit now two peaks and to contract somewhat in width, as shown in Figure 2(c).

In ice each proton is surrounded by several equally near magnetic neighbors. Also, a certain amount of local motion persists in ice well below the freezing point. This combination of more or less equidistant spacing of magnetic neighbors and of local molecular motion causes a further contraction of line width and a line of only one peak; however, this line is much lower and broader than that of protons in a water sample.

The data obtained from nuclear resonance absorption studies such as the above have yielded some information of the solid state not previously obtainable by other methods. For example, with the aid of certain advanced theory using data obtained by these methods

Consequently, higher than that of the Indian's median measurement. It has been shown statistically that the very majority of this group of Indian people measure the percentage of their body weight, as that of the Indian people, in the Indian people. It is shown that in the case of this measurement.

In the case of a single group of Indian people, the results of the measurements in different parts of the Indian people are shown, with some results from the Indian people. The results of the measurements in different parts of the Indian people are shown in a table, with the results from the Indian people. The results of the measurements in different parts of the Indian people are shown in a table, with the results from the Indian people.

In the case of a single group of Indian people, the results of the measurements in different parts of the Indian people are shown, with some results from the Indian people. The results of the measurements in different parts of the Indian people are shown in a table, with the results from the Indian people. The results of the measurements in different parts of the Indian people are shown in a table, with the results from the Indian people.

In the case of a single group of Indian people, the results of the measurements in different parts of the Indian people are shown, with some results from the Indian people. The results of the measurements in different parts of the Indian people are shown in a table, with the results from the Indian people. The results of the measurements in different parts of the Indian people are shown in a table, with the results from the Indian people.

The data obtained from the Indian people are shown in a table, with the results from the Indian people. The results of the measurements in different parts of the Indian people are shown in a table, with the results from the Indian people. The results of the measurements in different parts of the Indian people are shown in a table, with the results from the Indian people.

it has been possible to determine not only the directions of the H-H lines with respect to the axes of the gypsum crystal, but also, with a precision of about 1 per cent, the distance between the two protons in the HOH molecule.¹⁹ X-ray analysis does not yield this information.

(2) Method of Bloch

In the Bloch method for detection of nuclear magnetic resonance there is employed a nuclear induction transformer in which the establishment of nuclear resonance causes a change in coupling between the primary and secondary of the transformer and thus a change in the output voltage. This method of nuclear induction is based on the following principle.^{20,21}

20. Bloch, F. Phys. Rev. 70, 460 (1946).

21. Halliday, D. Introductory Nuclear Physics. New York: John Wiley & Sons, 1950, pp. 518-520.

If a sample of water or paraffin, say, is placed in a strong magnetic field H_0 there will be a small resultant magnetic moment due to the nuclear moments of the protons. The gross external effect results entirely from those few protons, about 7 in each 2,000,000 at a field strength of 10,000 gauss, that represent the difference in population between the two states available to the proton. If the strong magnetic field H_0 is in the z-direction and a weak r-f oscillating field of angular frequency ω is applied at right angles

It has been suggested that the Commission not only the responsibility of the law
should be placed on the shoulders of the government, but also that
a committee of inquiry be set up to investigate the various reasons for the
in the law. It is suggested that the Commission should also be
formed.

(2) The Commission

On the 10th of the month of January 1900 the Commission
has been formed. It is a body of five members, including the
President of the Commission, and a number of other members. The
President of the Commission is a member of the Commission, and
the members of the Commission are the members of the Commission.
The Commission is formed by the members of the Commission, and
the members of the Commission are the members of the Commission.
The Commission is formed by the members of the Commission, and
the members of the Commission are the members of the Commission.

20. The Commission is formed by the members of the Commission.

21. The Commission is formed by the members of the Commission.

A. The Commission is formed by the members of the Commission.

22. The Commission is formed by the members of the Commission.
The Commission is formed by the members of the Commission.
The Commission is formed by the members of the Commission.
The Commission is formed by the members of the Commission.
The Commission is formed by the members of the Commission.
The Commission is formed by the members of the Commission.
The Commission is formed by the members of the Commission.
The Commission is formed by the members of the Commission.
The Commission is formed by the members of the Commission.
The Commission is formed by the members of the Commission.

to H_0 in the direction of the x-axis with instantaneous magnitude given by

$$H_x = \hat{H} \cos \omega t,$$

then the combined fields H_0 and H_x will produce magnetic polarization in the sample. This polarization may be represented by the vector \underline{M} , the magnetic moment per unit volume. Bloch has shown²⁰ with the aid of certain simplifying assumptions that for a fixed H_0 and a fixed ω , (1) the magnitude of \underline{M} is constant, (2) \underline{M} rotates about H_0 at the angular rate ω , and (3) \underline{M} makes a fixed angle Θ with the H_0 axis. The three components of \underline{M} are

$$M_x = M \sin \Theta \cos \omega t,$$

$$M_y = M \sin \Theta \sin \omega t, \quad \text{and}$$

$$M_z = M \cos \Theta,$$

where $|\underline{M}| = M$ and the angle Θ is given by the relation

$$\tan \Theta = \frac{\hat{H}}{2(H_0 - H_{or})},$$

H_{or} being the value of H_0 at nuclear resonance. If $H_0 \gg H_{or}$, since \hat{H} is not large, Θ will also be fairly small. Under the condition of nuclear magnetic resonance, however, $H_0 = H_{or}$ and Θ becomes 90° . Consequently at resonance the components of \underline{M} become

$$M_x = M \cos \omega t,$$

$$M_y = M \sin \omega t, \quad \text{and}$$

$$M_z = 0.$$

From the last equation it is seen that $M_z = M \cos \Theta$, which is the

to H in the direction of the wave with instantaneous frequency
 given by

$$H_{\omega} = \hat{H} \cos \omega t$$

From the condition $H_{\omega} = \hat{H} \cos \omega t$ and $H_{\omega} = \hat{H} \cos \omega t$ we obtain
 that the wave is a plane wave. The polarization can be determined by the
 vector \hat{H} . The electric vector has only one component. From the above we
 obtain the direction of the wave vector \hat{k} for a plane
 wave and a plane wave (1) the direction of \hat{k} is constant. (2) \hat{k} is constant
 about \hat{k} at the origin and \hat{k} is constant about \hat{k} at the origin and \hat{k} is constant
 about \hat{k} at the origin and \hat{k} is constant about \hat{k} at the origin.

$$M_{\omega} = M \cos \omega t$$

$$M_{\omega} = M \cos \omega t$$

$$M_{\omega} = M \cos \omega t$$

and

where $|\hat{k}| = k$ and the angle θ is given by the relation

$$\frac{\hat{H}}{2(H_0 - H_0)} = \cos \theta$$

Using the value of \hat{k} as constant frequency. It is \hat{k} and \hat{k} is constant
 \hat{H} is not constant θ will also be constant. Under the condition
 of constant frequency, constant \hat{k} and θ is constant. The
 frequency of the wave is constant.

$$M_{\omega} = M \cos \omega t$$

$$M_{\omega} = M \cos \omega t$$

$$M_{\omega} = M \cos \omega t$$

From the last equation it is seen that $\hat{k} = \cos \theta$ and θ is constant.

time independent part of \underline{M} , vanishes at resonance. Since M_x and M_y do not vanish, however, \underline{M} is perpendicular to the direction of H_0 and rotates about it at a frequency which is equal to that of the Larmor precession. This state with $\theta = 90^\circ$ corresponds to equal populations being in each of the two energy levels so that no resultant steady moment exists. In the derivation of this Bloch assumed that the nuclei are entirely free with no spin-lattice interaction. Since spin-lattice interaction is not zero in practice, this equalization with $M_z = 0$ at resonance is not a completely accurate picture of the situation existing at that time.

At the condition of nuclear magnetic resonance the component of \underline{M} perpendicular to the direction of H_0 increases rather suddenly in magnitude. This component of \underline{M} rotates with the angular frequency ω . As a result, a variable flux links a pickup coil which has its axis in the y-direction. The resultant e.m.f. induced in the pickup coil is passed to a receiver and measured.

The operation of this type of nuclear resonance spectrometer has been summarized in the following manner by the developers of this technique.²²

22. Bloch, F., Hansen, W.W., and Packard, M. Phys. Rev. 70, 479 (1946).

"A spherical sample is immersed in a field $H_{dc} + H_{ac} \cos 377t$ produced by a magnet. The nuclear moments oriented by this field are caused to precess by a driving field H_x . The precessing moments

the following table is a summary of the results of the experiments conducted in the laboratory of the U. S. Bureau of Entomology and Plant Quarantine, Washington, D. C., during the summer of 1934.

induce voltages in a receiver coil. These voltages, which vary in amplitude at a 60 cycles per second rate because of the variations in field H_0 , are amplified along with stronger constant amplitude signals. The leakage and the varying voltages due to precessing nuclei are mixed in the detector, the output of which then contains pulsating unidirectional current, the steady component due to leakage and the variations corresponding to the desired signal. The d.c. is removed by blocking condensers in the amplifier which increases the signal voltages to a magnitude suitable for operation of the cathode ray tube."

One definitely advantageous feature of this method is that it affords the opportunity to determine the sign of the magnetic moment of a nucleus. This method is suited also for the study of line widths and shapes.

(3) Super-regenerative Method

In the super-regenerative method of detecting nuclear resonances a super-regenerative oscillator is used both to establish nuclear resonance and to detect the induced voltage from it. Several variations in experimental arrangements have been adopted by different investigators using this method. Roberts²³ has described one such

23. Roberts, A. Rev. Sci. Inst. 18, 845 (1947).

arrangement. The apparatus with which the writer obtained his

about 1000 ft. in a vertical well. These depths, which were in
agreement with the depths of the water in the
lake, were 1000 ft. and 1000 ft. respectively. The
depths of the water in the lake were 1000 ft. and 1000 ft.
respectively. The depths of the water in the lake were 1000 ft.
and 1000 ft. respectively. The depths of the water in the lake
were 1000 ft. and 1000 ft. respectively. The depths of the water
in the lake were 1000 ft. and 1000 ft. respectively. The depths
of the water in the lake were 1000 ft. and 1000 ft. respectively.

The depths of the water in the lake were 1000 ft. and 1000 ft.
respectively. The depths of the water in the lake were 1000 ft.
and 1000 ft. respectively. The depths of the water in the lake
were 1000 ft. and 1000 ft. respectively. The depths of the water
in the lake were 1000 ft. and 1000 ft. respectively. The depths
of the water in the lake were 1000 ft. and 1000 ft. respectively.

(1) Depth-temperature curves

The depth-temperature curves of the lake were obtained by means
of a depth-temperature recorder. The recorder was used to obtain
the depth-temperature curves of the lake. The recorder was used to
obtain the depth-temperature curves of the lake. The recorder was
used to obtain the depth-temperature curves of the lake. The recorder
was used to obtain the depth-temperature curves of the lake. The
recorder was used to obtain the depth-temperature curves of the lake.

The depth-temperature curves of the lake were obtained by means
of a depth-temperature recorder. The recorder was used to obtain
the depth-temperature curves of the lake. The recorder was used to
obtain the depth-temperature curves of the lake. The recorder was
used to obtain the depth-temperature curves of the lake. The recorder
was used to obtain the depth-temperature curves of the lake. The
recorder was used to obtain the depth-temperature curves of the lake.

The depth-temperature curves of the lake were obtained by means
of a depth-temperature recorder. The recorder was used to obtain
the depth-temperature curves of the lake. The recorder was used to
obtain the depth-temperature curves of the lake. The recorder was
used to obtain the depth-temperature curves of the lake. The recorder
was used to obtain the depth-temperature curves of the lake. The
recorder was used to obtain the depth-temperature curves of the lake.

measurements was similar in basic principle to that described by Roberts but differed in several details. The principle of operation to be described for the apparatus of this section will apply equally well to either arrangement.

The general layout of the apparatus used by the writer is exhibited schematically in the diagram of Figure 3. The oscillator shown in the diagram with connections to a coil in which a test tube has been inserted is actually placed in its entirety between the pole faces of the large magnet. This oscillator is of the super-regenerative type and from that fact this method derives its name.

In the super-regenerative method the triode of the super-regenerative oscillator produces self-sustained oscillations. For this, adequate feed-back from plate to grid is required to supply all the losses in the circuit. However, instead of allowing continued self-sustained oscillation, an externally generated modulating signal of appropriate strength called the quench voltage is applied either to the grid or to the plate of the triode. This sinusoidal quench voltage causes the overall gain of the triode to be varied in such a way that the self-sustained oscillation of the triode starts and stops once during each quench cycle.

Detection of nuclear magnetic resonance is accomplished in the following manner. When the quench voltage varies from a value which prevents oscillation to one which permits it to begin, the envelope of the current in the tank circuit begins to increase approximately exponentially and, if not inhibited, would reach a saturation value at which feed-back energy would just equal losses. The maximum

amplitude of oscillation reached and the integrated energy of the oscillation pulse are functions of the effective Q of the resonant circuit, the feed-back gain, and the quench amplitude and frequency.

When the oscillator is placed between the pole faces of a magnet and a sample containing the desired type of nucleus, say protons, is placed in the inductance coil of the tuned circuit, the nuclear spin axes of the protons will precess in the direction of the magnetic field at the Larmor frequency. When the magnetic field strength is of such value that the Larmor and the oscillator frequencies are the same, the r-f oscillation at the Larmor frequency will induce emission or absorption of radiation by the protons involving reorientation of their nuclear spin axes in the magnetic field. Since at thermal equilibrium in, say, a field of 10,000 gauss there are in each 2,000,000 hydrogen atoms some 7 more protons in the lower energy than in the higher energy nuclear spin state, there will be a net absorption of r-f radiation by the protons, the effective resistance of the tuned circuit will be increased and there will be a decrease in the rate at which oscillation builds up. Thus the integrated pulse will be less under conditions of resonance, a feature of the super-regenerative method which may be detected by appropriate methods to allow observation of the nuclear resonance.

The quench frequencies employed by the writer in his measurements were of the order of 2 to 10 kilocycles during the quench periods 100 to 500 microseconds. The thermal relaxation time for protons in water at room temperature has been measured²⁴ and found

The first thing I noticed when I stepped out of the car was the heat. It was a sticky, oppressive heat that seemed to wrap around me. I had heard that the weather in the South was terrible, but I didn't realize it would be this bad. I was used to the cool, crisp air of the North, and this was a shock. I took a deep breath and tried to ignore the heat. I had a long drive home, and I needed to get some rest. I pulled out of the parking lot and drove down the main road. The houses were small and close together, and the streets were narrow. I saw a few people walking on the sidewalks, but they didn't seem to mind the heat. I drove for about an hour and then I saw a sign that said "Welcome to the South". I smiled and continued on my way. The heat was still there, but I was starting to get used to it. I was in the South, and I was going to make the best of it.

24. Hahn, E.L. Phys. Rev. 76, 143 (1949)

to be 2.33 ± 0.07 seconds. Thus, in the case of the water sample, during that period of the quench cycle when the tuned circuit is permitted to oscillate there will be a very few more protons raised to the higher energy level than will be lowered to the level of lesser energy. During the quenched period there will be a tendency for thermal equilibrium to be re-established among nuclei, so when oscillation is again initiated more protons will again be elevated than lowered in energy state. This process gives a net absorption of energy from the super-regenerative oscillator for each cycle of the quench oscillations. It is this net absorption of energy which results in the Q of the circuit being lowered and the detection of resonance by the super-regenerative method being made possible.

Another feature of this method which should be noted is that the buildup of oscillation in the super-regenerative oscillator is exponential or faster and, if it starts from residual noise voltage in the tank circuit, may possibly cover a significant fraction or all of the quench cycle period without reaching saturation. If, however, when the quench cycle commences there is in the tuned circuit an external signal of the same frequency, the oscillation will build up from this external signal and either will reach a larger amplitude within the length of time permitted by the quench cycle, or else will reach saturation sooner. Although the external signal does not affect the rate of buildup of oscillation, the integrated pulse energy will nevertheless be greater. This can possibly be of

The first of these is the fact that the system is not a simple one. It is a complex system, and the complexity is not only in the number of components, but also in the way they are interconnected. The second is the fact that the system is not a static one. It is a dynamic system, and the dynamics are not only in the way the components interact, but also in the way the system evolves over time. The third is the fact that the system is not a linear one. It is a non-linear system, and the non-linearity is not only in the way the components interact, but also in the way the system evolves over time. The fourth is the fact that the system is not a deterministic one. It is a stochastic system, and the stochasticity is not only in the way the components interact, but also in the way the system evolves over time. The fifth is the fact that the system is not a single one. It is a multi-scale system, and the multi-scale nature is not only in the way the components interact, but also in the way the system evolves over time. The sixth is the fact that the system is not a single one. It is a multi-scale system, and the multi-scale nature is not only in the way the components interact, but also in the way the system evolves over time. The seventh is the fact that the system is not a single one. It is a multi-scale system, and the multi-scale nature is not only in the way the components interact, but also in the way the system evolves over time. The eighth is the fact that the system is not a single one. It is a multi-scale system, and the multi-scale nature is not only in the way the components interact, but also in the way the system evolves over time. The ninth is the fact that the system is not a single one. It is a multi-scale system, and the multi-scale nature is not only in the way the components interact, but also in the way the system evolves over time. The tenth is the fact that the system is not a single one. It is a multi-scale system, and the multi-scale nature is not only in the way the components interact, but also in the way the system evolves over time.

importance because of the following effect.

At the condition of resonance the nuclear spin axes of the sample precess about the direction of the magnetic field with the same frequency as the r-f oscillation of the tuned circuit. This precession continues during the period when the triode oscillation is quenched. When the quench voltage again reaches a value which allows oscillation to be resumed, it is possible that an e.m.f. may be induced in the oscillator coil as a result of persisting precession of the vector M as described in the Bloch method. If so induced this e.m.f. will alternate at the Larmor frequency which, at the resonance condition, would provide an external signal of the correct frequency to cause oscillation to start building up immediately in the tuned circuit. Being started immediately in this manner causes the oscillation to reach a larger amplitude or perhaps to reach saturation sooner than would be the case if the oscillation were built up from random noise. Thus in the event that the oscillation is built up from an induced e.m.f. due to nuclear precession, the integrated energy pulse of the oscillation will differ from that of non-resonance.

In either case, whether the Q of the circuit is changed at resonance or whether the precessing nuclei act like a signal generator in the tuned circuit of the oscillator, the integrated pulse of the r-f oscillation will be different from that of the non-resonance condition and thus may be detected and nuclear resonance observed. It is entirely possible that both effects may occur simultaneously in varying proportions under usual operating conditions.

The spectrum of a super-regenerative nuclear magnetic resonance

...months in order for us to understand the situation.

[illegible]

detector is not a single line spectrum but consists of a central resonance signal corresponding to the fundamental resonance frequency ν_0 and usually one or more pairs of sidebands. These sidebands occur at frequencies ν_s , given by

$$\nu_s = \nu_0 \pm n \nu_{qu},$$

in which ν_{qu} is the frequency of the a.c. quench voltage and assumes the values of successive small integers. Resonance absorption by the sample occurs and under proper test conditions may be detected as the magnetic field passes through that strength necessary to produce Larmor precession of the nuclei at the frequency ν_s for each of this limited number of sidebands.

The circuit of the super-regenerative oscillator used by the writer is shown in the diagram of Figure 4. In this circuit the quench frequency voltage is applied to the grid of the oscillator triode. Certain other investigators have applied the quench voltage to the plate. This last arrangement of the quench voltage at one time was used with the apparatus employed in this work but was abandoned because of the high noise to signal ratio resulting from this arrangement. An oscillator circuit of this latter type with quench voltage applied to the plate of the triode as used by Roberts²³ is shown in Figure 5.

(4) Heterodyne Method

In the heterodyne method a weakly oscillating detector is modulated by the nuclear absorption. Such a weakly oscillating detector

... of the ...

... of the ...

... of the ...

... of the ...

$$x^2 = y^2 + z^2$$

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...

... of the ...


... of the ...

... of the ...

(1) ...

... of the ...

... of the ...



39

CAPACITIES A AND B SELECTED
TO GIVE APPROPRIATE FREQUENCIES
FOR NUCLEI BEING COMPARED.
SAMPLE PLACED INSIDE COIL C.

Pole Face Super-Regenerative Oscillator

FIGURE 4

THE
SUPER-REGENERATIVE
RESONANCE DETECTOR
(AFTER ROBERTS)^{2,3}




FIGURE 5.



is called an autodyne; in this case the signal from the autodyne beats against an incoming c-w signal to produce a heterodyne signal. A frequently difficult but required condition is that the oscillation of the autodyne be extremely weak, for in this method the strength of the resonance signal is found to diminish as that of the autodyne signal increases, apparently because of saturation of the nuclear absorption. This saturation evidently results from the fact that the total power which can be absorbed by a sample small enough to allow suitable homogeneity of the magnetic field is fairly minute. For N atoms with spin I at temperature T whose Larmor frequency is ν and relaxation time T_1 , the power which can be absorbed is

$$P = \frac{2I}{I+1} \cdot \frac{N}{T_1} \cdot \frac{(h\nu)^2}{kT}.$$

For a sample of 1 c.c. of water at 300°K with ν_0 at 30 mc/sec, P will be approximately 10^{-9} watts.²³

The super-regenerative circuit shown in Figure 5 can be adapted for use as an autodyne detector if the quench voltage supply is removed and the plate voltage properly adjusted. The circuit shown in this figure would probably not be as sensitive as a super-regenerative oscillator.²³ However, improved and more sensitive forms of this method have been devised.²⁵

25. Pound, R.V. Phys. Rev. 72, 527 (1947).

Nuclear magnetic resonance may be observed in either of two

It is noted on subject's file that the subject has been in contact with the following persons:

[illegible]

$$P = \frac{I_2}{1 + I_2} \cdot \frac{V}{T} \cdot \frac{A_2(A_1)}{T_2}$$

9. The number of the page is 100, and the number of the page is 100.

The most representative example of the (1) type is the
the use of an arbitrary interval 17 in figure 10. The
covered and the fixed voltage source. The circuit shown
in this figure would provide an example of a non-linear
five amplifier.¹² However, improved and more sensitive forms of
this method have been devised.¹³

28. Brown, R. V., *et al.*, 1995. *Ann. Entomol. Soc. Am.* 88: 100-105.

© 2002 by Blackwell Publishers Ltd. *Journal of Internal Medicine* 252: 429–436

ways when using the autodyne detector. If the output is viewed by an a-m receiver the absorption curve for a liquid sample appears much the same as that of the absorption line in Figure 2 (a). If f-m detection is used, the signal has the form of an anomalous dispersion curve. This f-m effect results from the oscillator being slightly detuned by nuclear resonance effects.

page when using the ordinary battery. In the course of time it
is also possible the battery may be a little more
than the time it takes to be charged. It is also possible
the battery is used, but it may be the fact of an electrical
discharge may. This is why people have the battery
being already damaged by earlier excessive use.

III. EXPERIMENTAL ARRANGEMENTS

(a) Apparatus

The experimental arrangement used in conducting the tests described in this report is indicated in the block diagram of Figure 3. The magnet used to produce the strong magnetic field was a double yoke type with high voltage, low current windings and an adjustable air gap. The gap width used in these experiments was approximately $3/4$ ". Pole pieces of $3\frac{1}{2}$ " diameter were used in early work, but all measurements included in this report were made with 6" pole pieces because of the better magnetic field homogeneity obtainable with the larger diameter pole faces. The magnet current was supplied by an electronically controlled current regulator with a clockdrive mechanism operating a potentiometer in the control circuit. This current regulator was capable of increasing or decreasing the magnetic field at a rate of about 2 gauss per minute over a field strength range of some 2,000 to 10,000 gauss. Except under unfavorable conditions seeming usually to stem at irregular hours from rapid uneven variations in the 110 volt 60 cycle a.c. power supplied to the laboratory, the current regulator was capable of maintaining the magnet field current constant to about one part in thirty thousand.

The strong magnet field H_0 was modulated by a 45 cycles per second current applied to small field modulating coils called "sweep coils," one being clamped on each pole piece. The 45 cycle current for these windings was supplied by a signal from an audio frequency signal generator amplified by a push-pull amplifier. This small

alternating modulation H_m to the strong field H_0 caused the resonance signal to be repeated periodically at any time when the resonance frequency happened to be within the limits corresponding to the field range $H_0 \pm H_m$. The resonance signal was observed usually on an oscilloscope with its sweep synchronized to the signal generator producing the 45 c.p.s. field sweep signal and was recorded by an Esterline-Angus recorder.

The samples used in these experiments were either saturated solutions or concentrated acids. Approximately one c.c. of the sample was placed in a $\frac{1}{8}$ " diameter test tube which in turn was placed in the inductance coil of the super-regenerative oscillator tank circuit. The oscillator was so located in the magnet gap that the sample would occupy that portion of the field found by careful testing to be most homogeneous.

The tank circuit of the oscillator was tuned to a frequency somewhere in the range between 1 to 10 megacycles per second, depending upon the nuclear g-factor of the unknown reported by molecular beam experimenters and the magnetic field strength desired to be used for the experiment. This frequency could, after once being set by appropriate choice of fixed capacitance condensers, be varied within narrow limits by means of a small variable condenser in the tank circuit. In most cases the small variation in oscillator frequency permitted by the tuning condenser would not cover the frequency range required for comparison of the resonance frequency for the sample of undetermined g-factor with that of the known sample. This consideration required that a second set of fixed capacitance condensers be installed in the

pole face oscillator with necessary switching arrangements to permit rapid changing between the two frequency ranges.

The alternating quench voltage applied to the grid of the oscillator triode was variously set from 2.0 to 9.0 kilocycles and from 1.5 to 3.0 volts. These and other parameters were varied both in attempts to locate the resonance signal and, after the signal was found, to obtain a sharp, accurately measurable signal pattern. Other parameters varied for the same purpose were the magnetic field modulation strength, oscillator plate voltage, and the concentration and type of chemical compound used for the sample.

The signal from the pole face oscillator was sent through a narrow band amplifier tuned to filter out all alternating current components except the 45 cycles per second of the magnetic field modulation. The output of the narrow band amplifier was sent both to an oscilloscope and to a phase-detector. The output of this latter instrument was in turn passed to an Esterline-Angus recorder.

At resonance, corresponding to the condition

$$h\nu_0 = g_z \mu_m H,$$

the amplitude of oscillation of the super-regenerative oscillator is altered because of the previously discussed resonance phenomena. This amplitude change was observed visually on the oscilloscope trace and the time derivative of this change in amplitude of oscillation was recorded by the Esterline-Angus recorder from the output of the phase detector.

After resonance was observed and operating parameters were properly adjusted to obtain a signal suitable for measuring, the clock-

drive mechanism operating the potentiometer of the magnet current regulator was stopped and the field current regulator was allowed to maintain a constant current to the magnet field coils. The center of the resonance spectrum of the sample to be measured was accurately located by tuning the variable condenser in the pole face oscillator circuit and then measuring the oscillator frequency using the heterodyne frequency meter shown in Figure 3. With the magnet current still maintained constant, a sample of known value of nuclear g-factor was substituted for the unknown in the pole face oscillator and the oscillator frequency varied until resonance was obtained for this standard sample. With g_x and g_s denoting respectively the nuclear g-factors of the nucleus being studied and the nucleus used for standard, and ν_x and ν_s denoting their frequencies in the same magnetic field, g_x was obtained from the relation

$$g_x = g_s \frac{\nu_x}{\nu_s}$$

Several pairs of such readings commonly were obtained and the average of their ratios was taken as the value for the measurement. Probable errors of measurements were computed according to customary procedures.

(b) Modulation Improvements

The magnetic field modulation for the initial period of this report was achieved by means of only a single sweep coil clamped on one pole piece. It was wound with about 2500 turns of 32 gauge wire and was supplied with 45 cycle voltage produced by an audio signal generator and amplifier. Sweep coil voltage was varied ordinarily in the range from 20 to 60 volts as necessary to obtain optimum

...the ... of the ...
...the ... of the ...
...the ... of the ...

...the ... of the ...
...the ... of the ...
...the ... of the ...

...the ... of the ...
...the ... of the ...
...the ... of the ...

...the ... of the ...
...the ... of the ...
...the ... of the ...

...the ... of the ...
...the ... of the ...
...the ... of the ...

...the ... of the ...
...the ... of the ...
...the ... of the ...

...the ... of the ...
...the ... of the ...
...the ... of the ...

resonance signals.


After the measurements on Indium¹¹³ were completed, tests were made using a small induction coil and an a.c. voltmeter to ascertain whether field modulation was appreciably constant over the region occupied by the sample in the gap between the pole faces. Although the small coil had a width equal to almost half of the pole face gap, a decrease in induced voltage of approximately fifty per cent was noted when the coil was held against the center of the face of the pole piece on which the sweep coil was mounted and from there moved parallel with the magnet axis to a position against the opposite pole face.

As a result of these tests it was decided to mount a similar coil on the second pole piece and to construct a push-pull amplifier to provide the sweep field power for both coils. The circuit diagram of this amplifier is shown in Figure 6. The new coil was over-wound by a few hundred turns to allow margin for proper balancing of the two coils. After the new amplifier and sweep coil were installed and the sweep field balanced by removing a certain number of excess turns from the newer coil, the variation in voltage induced in the probe coil was reduced to five per cent or less.

(c) Homogenizing of the Magnetic Field

The preparatory work attempted by the writer prior to making any actual measurements was performed using $3\frac{1}{2}$ " diameter pole pieces. Prior to that it had been the custom to fill the test tubes with sample solutions to a height of about two and one-half

22.5-
22.5



Push-Pull Sweep Amplifier




FIGURE 6.

inches. Utilizing a relatively large amount of sample in this manner ordinarily gave a strong resonance signal, but the center of the resonance pattern usually was lacking in the degree of sharpness desired for maximum accuracy of measurement.

It appeared possible that lack of desirably sharp resonance signals might be attributable in part to inhomogeneity of the magnetic field over the volume occupied by the sample. To check this assumption, test runs were made using smaller amounts of the sample solution. This procedure immediately produced sharper resonance patterns. After a number of tests it was ascertained that acceptable resonance signal strengths usually could be obtained using a depth of liquid in the tube ranging from 1.0 to 1.5 cm. Appreciably less than this amount for many nuclides frequently gave an undesirably weak signal. Exceptions to this last statement were the proton and the deuteron, both of which were found to give strong signals from only a few drops of H_2O or D_2O , respectively.

In a further attempt to achieve better field homogeneity, the $3\frac{1}{2}$ " pole pieces were replaced by others of 6" diameter. This change also produced an improvement in the sharpness of the resonance pattern but carried with it the slight disadvantage of placing a lower limit on the maximum field strength obtainable. This limitation was inconvenient upon occasions but was never found to be unacceptably restrictive.

In spite of the fact that the 6" pole pieces gave better resonance patterns it appeared probable that an additional improvement in field homogeneity could be obtained by removing a few slight burrs from the

pole faces. Further investigation into possible causes of field inhomogeneity also revealed that neither pole face was acceptably flat nor were the two faces sufficiently parallel. To correct all except the last of these deficiencies each pole face was carefully re-machined to make it as flat as possible and then was ground and polished by hand using emery dust and jeweler's rouge as grinding compounds. These efforts eventually produced reasonably flat and smooth pole face surfaces.

After the pole pieces were reinstalled a test was made to determine the most homogeneous part of the magnetic field for sample occupancy. For this check a small probe oscillator was employed using a sample of only some five or six drops of water from which to obtain proton resonance. Initial results indicated that the magnetic field even near the center of the pole faces varied as much or more than five gauss when the proton probe was moved a half inch in practically any direction. The pattern of variation indicated however, that shimming one side of one of the pole pieces might produce beneficial results. A long process of trial and error shimming using thin non-magnetic shims eventually produced a field which varied appreciably less than one gauss over a region sufficiently large to accommodate the desired height of liquid in a $\frac{1}{8}$ " diameter test tube.

The deviation of the magnetic field from true homogeneity both before and after the remedial efforts described above is indicated in the graphs of Figure 7.

both cases. Further investigation has revealed, however, that this

relationship also extends to the other two cases.

That we have the same relationship holds. It follows that

these two cases are also related to the first case.

It follows, therefore, that the relationship between the

three cases is a general one, and that the relationship

between the first two cases is a special case of the

general one.

After the first case has been established, it is not

difficult to see how the second case is a special case of the

general one. For this case is a special case of the

general one, and the first case is a special case of the

general one. It follows, therefore, that the

relationship between the first two cases is a special case of the

general one. It follows, therefore, that the

relationship between the first two cases is a special case of the

general one. It follows, therefore, that the

relationship between the first two cases is a special case of the

general one. It follows, therefore, that the

relationship between the first two cases is a special case of the

general one. It follows, therefore, that the

relationship between the first two cases is a special case of the



general one. It follows, therefore, that the

relationship between the first two cases is a special case of the

general one. It follows, therefore, that the

relationship between the first two cases is a special case of the


general one. It follows, therefore, that the



DISTANCE (INCHES) FROM CENTER OF WEST POLE FACE

RESULTS OF HOMOGENIZING THE MAGNETIC FIELD

FIGURE 7.



(d) Results of Improvements

The results obtained from efforts spent to homogenize the magnetic field are well illustrated by differences appearing in the "before and after" resonance signals traced by the Esterline-Angus recorder. The typical "before" trace of a sharp resonance pattern appeared as an uninterrupted series of slow swings of the recorder pen in alternate directions. Pronounced hesitations in the movements of the pen occurred in this type of pattern only at each side limit of the recorder paper when the amplitude of the resonance signal exceeded the maximum capability of the recorder. A typical pattern of this type obtained from Indium¹¹⁵ is shown in Figure 8. In this pattern one might estimate that there are three resonance signals corresponding to the central signal and one pair of side-band resonances. Even so, there is nothing in the pattern which can be chosen as serving to separate the three theoretically distinct resonance signals.

The series of resonance signals in this pattern would be seen on the oscilloscope in the following sequence. Upon reaching the first resonance band the small fluctuations of the oscilloscope caused by random noise would be replaced by a steady buildup to a sustained peak of large amplitude followed presently by a steady decrease to a sustained valley of similar amplitude. This peak-valley would be twice repeated by the signal recorded in Figure 8. The trace on the oscilloscope in moving from valley to peak and vice-versa would do so without any significant hesitation as the trace crossed the horizontal axis. At the conclusion of the signal

The results obtained from these tests are presented in the

table which will illustrate the differences appearing in the
 "before and after" treatment of the material.
 The "before" test is a very common type
 known as an unbalanced test in that only one of the
 two treatments is compared. The results are the same
 as the results obtained in this type of test only in that
 the results are not the results of the treatment
 but the results of the treatment of the material. A typical
 result of this type of test is shown in Figure 2.
 In this figure the results show that the results
 obtained are the results of the treatment of the material
 and not the results of the treatment of the material.
 The results are the results of the treatment of the material
 and not the results of the treatment of the material.
 The results are the results of the treatment of the material
 and not the results of the treatment of the material.

The results of these tests are presented in the
 table which will illustrate the differences appearing in the
 "before and after" treatment of the material.
 The "before" test is a very common type
 known as an unbalanced test in that only one of the
 two treatments is compared. The results are the same
 as the results obtained in this type of test only in that
 the results are not the results of the treatment
 but the results of the treatment of the material. A typical
 result of this type of test is shown in Figure 2.
 In this figure the results show that the results
 obtained are the results of the treatment of the material
 and not the results of the treatment of the material.
 The results are the results of the treatment of the material
 and not the results of the treatment of the material.
 The results are the results of the treatment of the material
 and not the results of the treatment of the material.




FIGURE 8.

the oscilloscope would again display random noise effects.

A different resonance pattern representative of the sharp "after" type is shown in Figure 9. In this pattern obtained from Chlorine³⁷ it may be seen that the central resonance signal is clearly separated from each of the two signals of the one sideband pair which are in evidence. This separation appeared on the oscilloscope as a return of random noise fluctuations for a period of approximately one and a half minutes between the separate resonance signals. It may also be noticed that the width of each signal is narrower, a feature surely attributable to better field homogeneity.

Another new feature which has not yet been adequately explained is the following phenomenon to be seen in Figure 9. Observing the direction of the initial excursions of the recorder pen in each of the three separate resonance signals one notices in the right hand signal that the first excursion is toward the top of the paper; that of the central signal is toward the bottom, while that of the left-hand signal is again upward. This alternation of direction of initial swings of the recorder pen is not confined merely to first order sidebands but has been noticed in all side band resonance signals observed since the field homogeneity was improved.

It will be recalled that the Esterline-Angus recorder receives its input from the super-regenerative oscillator through a narrow band amplifier and a phase detector. No conclusion has yet been reached why the resonance signal as detected by this arrangement should behave in the above manner. Studies, however, are being

The conditions would again likely be met.

A different parameter (being representative of the data

*which is shown in Figure 2. In this figure, the data

showed that it may be seen that the central parameter is

slightly increased from each of the two sides of the two

part with the in between. This operation appears to be

different as a result of the two sides of the two sides of

operation, and a half of the data between the two sides

operation. It may be noted that the data of each side is

operation, a further half of the data is shown in the

operation.

Figure 2 shows that the data is not yet been completely

in the data, the data is shown in Figure 2. The data

operation of the data, the data is shown in Figure 2. The data

the data, the data is shown in Figure 2. The data

operation, the data is shown in Figure 2. The data

of the data, the data is shown in Figure 2. The data

data, the data is shown in Figure 2. The data

operation, the data is shown in Figure 2. The data

operation, the data is shown in Figure 2. The data

operation, the data is shown in Figure 2. The data

It may be noted that the data is shown in Figure 2. The data

the data, the data is shown in Figure 2. The data

data, the data is shown in Figure 2. The data

operation, the data is shown in Figure 2. The data

operation, the data is shown in Figure 2. The data




FIGURE 9.

continued by other personnel to attempt to explain this phenomenon.

A definite advantage in one phase of making frequency measurements on resonance signals accrued from the work on the pole pieces. In the older type of pattern one was forced to choose the point in the pattern representing the center of symmetry. It can be seen in Figure 8 that only the point indicated by X satisfies this condition. For an accurate measurement to be obtained it was essential that the swing of the pen across the region of the paper near the X occur without any hesitation. Unfortunately, in practice this condition is difficult to obtain.

With the newer resonance pattern in which each side band resonance signal was clearly distinguishable it was discovered to be possible to shift to a sharp side band if the central resonance signal happened not to be sharp. Having shifted to a side band to obtain frequency measurements one then only had to apply the correction

$$n \nu_{gu}$$

to the observed side band resonance frequency ν_s to obtain the desired but unobserved central resonance frequency ν_0 . The relation employed for this purpose is

$$\nu_s = \nu_0 + n \nu_{gu},$$

in which n , the order of the side band, is given the appropriate algebraic sign.

Measurements obtained from side bands were in no manner less accurate than those obtained from the central resonance signal.

Although most nuclei seldom exhibited more than two or three pairs of side bands in a resonance pattern, at least ten such pairs

of side bands have been observed frequently in resonance signals produced by the deuteron in the homogenized magnetic field. The upper limit on n for the deuteron can not be stated with accuracy because, in the interest of economy of time and ease of measurement, efforts were usually exerted to limit the number of side bands in a pattern consistent with obtaining the desired degree of sharpness in the signal.

$$\omega_p = \gamma \omega^2 \mu = \gamma \mu \left(\frac{\omega}{\gamma} \right) \left(\frac{\gamma}{\omega} \right) \omega$$

The value of the ratio (ω_p/ω) was then calculated from the results obtained by using the above formula. The value used for the γ factor of the deuteron was

$$\gamma = 5.27982 \pm 0.0001$$

obtained from the value $(\gamma_p = 1.75946 \pm 0.00005) \mu_{\text{proton}}$

The probable error indicated on this value for the ratio

(ω_p/ω) is a measure of the fractional uncertainty of the value. Since the existence of systematic errors as large as the ratio of the above formula can not be excluded, the probable error of the above value has been calculated using the systematic error of that ratio as the probable error of the value

$$(\omega_p/\omega)$$

The probable error of the above value is approximately 0.0001. The systematic error of the above value is approximately 0.0001. The total error of the above value is approximately 0.0001. It is therefore not possible to state the value of the ratio with greater accuracy.

of this kind have been observed in various places
and by the authors in the same localities. The
effect of the light on the color of the
eye, however, in the absence of any other cause of color-
change, appears to be directly related to the nature of the light
in a, the wavelength of which is the same as that of the
light in the eye.

The effect of the light on the color of the
eye, however, in the absence of any other cause of color-
change, appears to be directly related to the nature of the light
in a, the wavelength of which is the same as that of the
light in the eye. The effect of the light on the color of the
eye, however, in the absence of any other cause of color-
change, appears to be directly related to the nature of the light
in a, the wavelength of which is the same as that of the
light in the eye.

The effect of the light on the color of the
eye, however, in the absence of any other cause of color-
change, appears to be directly related to the nature of the light
in a, the wavelength of which is the same as that of the
light in the eye.

The effect of the light on the color of the
eye, however, in the absence of any other cause of color-
change, appears to be directly related to the nature of the light
in a, the wavelength of which is the same as that of the
light in the eye.

The effect of the light on the color of the
eye, however, in the absence of any other cause of color-
change, appears to be directly related to the nature of the light
in a, the wavelength of which is the same as that of the
light in the eye.

IV. EXPERIMENTAL RESULTS

The measurements to be described in this section were obtained by comparing in the same magnetic field the resonance frequency of each unknown with that of a known sample used as a standard. The magnetic moment of the unknown sample was then obtained from the relation

$$\mu_x = g_x I_x = g_p \left(\frac{\nu_x}{\nu_s} \right) \left(\frac{\nu_s}{\nu_p} \right) I_x$$

In each case the ratio ($\nu_{\text{standard}} / \nu_{\text{proton}}$) has been obtained from recent results published by other investigators. The value used for the g-factor of the proton was

$$g_p = 5.58536 \pm 0.00012,$$

obtained from the value $\mu_p = (2.79268 \pm 0.00006) \mu_n$.

The probable error indicated in each table for the ratio ($\nu_{\text{unknown}} / \nu_{\text{standard}}$) is a measure of the internal consistency of the data. Since the existence of systematic errors as large as one part in twenty thousand can not be excluded, the probable error of the magnetic moment has been computed using the systematic error if that error exceeded the probable error of the ratio ($\nu_{\text{unknown}} / \nu_{\text{standard}}$).

The results here have not been corrected for diamagnetic effects of the extra-nuclear electrons nor for the slight paramagnetic effect of a magnetic catalyst in those cases in which it is indicated that such a catalyst was used.

The measurements as described in this section were obtained by averaging in the same manner the two series of measurements of the same sample and of a second sample. The relative error of the measurements was then calculated from the relation

$$I_{\text{rel}} = \frac{I_1}{I_2} = \frac{I_1}{I_2} \left(\frac{I_1}{I_2} \right) \left(\frac{I_1}{I_2} \right) \left(\frac{I_1}{I_2} \right)$$

In each case the value of I_{rel} has been calculated from the results published by other investigators. The value used for the calculation of the error was

$$I_{\text{rel}} = 2.5 \times 10^{-3}$$

$$I_{\text{rel}} = 2.5 \times 10^{-3} \pm 0.0001$$

The results were calculated in each case from the value

(1) $I_{\text{rel}} = 2.5 \times 10^{-3}$ is a measure of the relative sensitivity of the detector. Since the calculation of I_{rel} is based on the assumption that the detector is not saturated, the relative error of the results would be increased if the detector were saturated. The results would be increased if the detector were saturated.

$$I_{\text{rel}} = 2.5 \times 10^{-3}$$

The results have been calculated for the detector of the type described above. The results are given in Table I. A comparison of the results with those obtained by other investigators is given in Table II. The results are given in Table III. A comparison of the results with those obtained by other investigators is given in Table IV.

(a) Indium¹¹⁵

The measurement of the nuclear magnetic moment of In¹¹⁵ was obtained from one c.c. of a saturated solution of In(NO₃)₃ in 30% HNO₃ to which had been added a small amount of Mn(NO₃)₂ as a catalyst. The standard used for comparison was Sc⁴⁵ in one c.c. of a saturated solution of ScCl₃. The value for the ratio

$$\frac{\nu_{\text{In}^{115}}}{\nu_{\text{Sc}^{45}}} = 0.242939 \pm 0.000003$$

is that due to Huxton.²⁶ The data pertaining to this measurement

26. Huxton, D.M. Phys. Rev. **78**, 806 (1950).

Table 1.
Data for Indium¹¹⁵

Unknown Nuclide	Indium ¹¹⁵
Standard Nuclide	Scandium ⁴⁵
$\frac{\nu_{\text{In}^{115}}}{\nu_{\text{Sc}^{45}}}$	0.902292 ± 0.000005
$\frac{\nu_{\text{Sc}^{45}}}{\nu_{\text{H}}}$	0.242939 ± 0.000003
$\frac{\nu_{\text{In}^{115}}}{\nu_{\text{H}}}$	0.219202 ± 0.000012
$g_{\text{I}}(\text{In}^{115})$	1.22432 ± 0.00012
$\mu_{\text{I}}(\text{In}^{115})$	(5.50945 ± 0.00015)

are summarized in Table 1.

(b) Chlorine³⁷

The measurement of the nuclear magnetic moment of Cl³⁷ presented in Table 2 was obtained using a saturated solution of approximately one c.c. of LiCl to which had been added a small amount of MnCl₂ as a magnetic catalyst. The deuteron resonance frequency used as a standard was obtained from a one c.c. sample of D₂O. The value of the ratio

Table 2.

Data for Chlorine³⁷

Unknown Nuclide	Chlorine ³⁷
Standard Nuclide	Deuteron
$\frac{\nu_{Cl^{37}}}{\nu_{D^2}}$	0.531632 ± 0.000036
$\frac{\nu_{D^2}}{\nu_{H^1}}$	0.153506
$\frac{\nu_{Cl^{37}}}{\nu_{H^1}}$	0.0816027 ± 0.0000054
$E_I(Cl^{37})$	0.455848 ± 0.000032
$\mu_I(Cl^{37})$	0.683722 ± 0.000048

$\frac{\nu_{D^2}}{\nu_{H^1}} = 0.153506$ is taken as being accurate to six significant figures.²⁷

27. Smaller, B., Yasaitis, E., and Anderson, H.L. Phys. Rev. **80**, 137(A) (1950).

(c) Nitrogen¹⁴

The measurement for N¹⁴ summarized in Table 3 was obtained using approximately one c.c. of concentrated HNO₃. Rb⁸⁵ in a one c.c. saturated solution of RbCl was used as the standard. The ratio ($\nu_{Rb^{85}}/\nu_{H^1}$) was obtained from results by Yasaitis and Smaller in which the ratio was given as²⁸

$$\frac{\nu_{H^1}}{\nu_{Rb^{85}}} = 10.357105 \pm 0.00001.$$

28. Yasaitis, E., and Smaller, B. Phys. Rev. 82, 750 (1951).

Table 3.

Data for Nitrogen¹⁴

Unknown Nuclide	Nitrogen ¹⁴
Standard Nuclide	Rubidium ⁸⁵
$\frac{\nu_{N^{14}}}{\nu_{Rb^{85}}}$	0.748425 \pm 0.000019
$\frac{\nu_{Rb^{85}}}{\nu_{H^1}}$	0.096552 \pm 0.0000001
$\frac{\nu_{N^{14}}}{\nu_{H^1}}$	0.072262 \pm 0.000002
$\epsilon_I(N^{14})$	0.40361 \pm 0.00002
$\mu_I(N^{14})$	0.40361 \pm 0.00002

$$P_{\text{measured}}(0)$$

The measurement of the $P_{\text{measured}}(0)$ is obtained by using a series of measurements of the $P_{\text{measured}}(0)$ at different values of the parameter α . The results are shown in Table 1. The values of $P_{\text{measured}}(0)$ are obtained by using the values of $P_{\text{measured}}(0)$ at different values of α and the values of $P_{\text{measured}}(0)$ at different values of α are shown in Table 1.

$$10.32102 \pm 0.0001 = \frac{1.4}{1.4}$$

Table 1. Values of $P_{\text{measured}}(0)$ at different values of α .

α	$P_{\text{measured}}(0)$	$P_{\text{measured}}(0)$
0.0000	1.4000	1.4000
0.0001	1.4000	1.4000
0.0002	1.4000	1.4000
0.0003	1.4000	1.4000
0.0004	1.4000	1.4000
0.0005	1.4000	1.4000
0.0006	1.4000	1.4000
0.0007	1.4000	1.4000
0.0008	1.4000	1.4000
0.0009	1.4000	1.4000

V. DISCUSSION OF RESULTS

During the time which has elapsed since the measurements listed in Section IV. were obtained, another member of the laboratory, Dr. Yu Ting, has obtained measurements of the nuclear magnetic moments of both Cl^{35} and In^{113} . In the measurement of Cl^{35} the standard used was Cl^{37} in the same sample. In the case of In^{113} , In^{115} was used as a standard. The In^{113} was contained in enriched proportions in a sample supplied on loan by the U.S. Atomic Energy Commission. The results tabulated in Tables 4 and 5 include the results obtained both by Dr. Ting and by the writer. In addition there are listed in Table 4 for the purpose of later discussion the experimental values of the quadrupole moments for the various nuclides as well as the nuclear magnetic moments of In^{113} and In^{115} obtained by the atomic beam method.

It will be noticed in Table 4 that the atomic beam values of the magnetic moments of both isotopes of Indium are less than the values obtained by magnetic resonance methods. These differences are much too large to be accounted for within the probable errors of the experiments. Similar discrepancies have been noted^{29,30,31} for

29. Taub, H., and Kusch, P. Phys. Rev. **75**, 1481 (1949).

30. Pound, R.V. Phys. Rev. **73**, 1112 (1948).

31. Kusch, P. Phys. Rev. **73**, 584 (1948).

the isotopes Ga^{69} and Ga^{71} .

Table 4.

Data on μ_I^* and Q

Nuclide	Z	A-Z	I	μ_I (Mag. Res.)	μ_I (Atomic Beam)	Q (10^{-24} cm ²)
Cl ³⁵	17	18	3/2	0.820896 μ_n		-0.0795 \pm 5 a.
Cl ³⁷	17	20	3/2	0.683722		-0.0621 \pm 5 a.
In ¹¹³	49	64	9/2	5.49507	5.486 a.	1.144 b.
In ¹¹⁵	49	66	9/2	5.50945	5.500 a.	1.161 b.
N ¹⁴	7	7	1	0.40361		0.02 a.

*Values for μ_I by magnetic resonance method were obtained at this institution.

a. Mack, J.E. Revs. Mod. Phys. 22, 64 (1950).

b. Mann, A.K., and Kusch, P. Phys. Rev. 77, 427 (1950).

Table 5.

Comparison of Results with Those Obtained Elsewhere

Nuclide	μ_I (Mag. Res. from Table 4.)	μ_I (Proctor and Yu ³²)
Cl ³⁵	(0.820896 \pm 0.000050) μ_n	(0.8211 \pm 0.0001) μ_n
Cl ³⁷	0.683722 \pm 0.000048	0.6835 \pm 0.0001
In ¹¹³	5.49507 \pm 0.00020	5.4972 \pm 0.0010
In ¹¹⁵	5.50945 \pm 0.00015	5.5088 \pm 0.0010
N ¹⁴	0.40361 \pm 0.00002	0.40369 \pm 0.00006

32. Proctor, W.G., and Yu, F.C. Phys. Rev. 81, 20 (1951).

Table 1.

Mean and μ_2 and σ

Variable	1	2	3	4	5	6	7	8	9	10
μ_1	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
μ_2	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
σ	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

Values for μ_1 for complete response: values were obtained at this time.

1. μ_1 (1971) 2. μ_2 (1971) 3. μ_3 (1971) 4. μ_4 (1971) 5. μ_5 (1971)

6. μ_6 (1971) 7. μ_7 (1971) 8. μ_8 (1971) 9. μ_9 (1971) 10. μ_{10} (1971)

Table 2.

Regression of μ_1 on μ_2 and μ_3

μ_1 (1971) μ_2 (1971) μ_3 (1971)

from Table 1.

μ_1	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
μ_2	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
μ_3	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
μ_4	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
μ_5	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000

2. μ_1 (1971) 3. μ_2 (1971) 4. μ_3 (1971) 5. μ_4 (1971) 6. μ_5 (1971)

Foley³³ has investigated a suggestion by Townes that partial

33. Foley, H.M. Phys. Rev. 80, 288 (1950).

decoupling of the L and S vectors in the $^2P_{1/2}$ state in the applied magnetic field of the atomic beam apparatus might affect the hyperfine interaction in such a way that the nuclear moment appears to be altered. The results obtained by Foley and his co-workers using a non-relativistic one-electron wave function in developing an expression for the energy perturbation of the magnetic levels of the $^2P_{1/2}$ state have indicated that the effect does not quite account for the entire discrepancy observed. Foley has further pointed out that the ground states of such atoms as gallium, indium and thallium are subject to configuration interaction mixing with some excited states of the outer three electrons, which affects strongly the hyperfine splittings of the $^2P_{3/2}$ states. This fact might possibly account for the remaining discrepancy.

The values of the results obtained by the writer and also those by Dr. Ting as listed in Table 5 are seen to agree with those obtained by Proctor and Yu at Stanford within one digit at the fourth significant figure within the limits of probable errors. Results obtained by other investigators generally come within limits as close or smaller than this.

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

THE UNIVERSITY OF CHICAGO

(a) Shell Theory

The statement was made in Section I that no theory at present is capable of predicting magnetic moments of nuclides having non-zero spin. There is a model of the nucleus due to Schmidt,^{34,35,36} however,

34. Schmidt, T. Zeits. f. Physik 106, 358 (1937).

35. Fermi, E. Nuclear Physics. Chicago: The University of Chicago Press, 1950, p. 14.

36. Blatt, J.M., and Weisskopf, V.F. Theoretical Nuclear Physics. New York: John Wiley & Sons, 1952, pp. 38-52, 767-774.

which attempts to explain nuclear magnetic moments by assuming that the nuclear spin I in an odd-even or even-odd nucleus results from the motion of the one unpaired nucleon. The sum of the orbital and spin angular momenta of all other nuclear particles is assumed to be zero. For this model we have

$$\begin{aligned}\underline{I} &= \underline{L}_K + \underline{S}_K && \text{AND} \\ \underline{\mu}_I &= \frac{e\hbar}{4\pi Mc} (g_L \underline{L}_K + g_S \underline{S}_K),\end{aligned}$$

where g_L and g_S are gyromagnetic ratios with values

$$g_L = 1 \quad \text{and}$$

$$g_S = g_p = 1$$

if the odd particle is a proton, or

$$g_L = 0 \quad \text{and}$$

$$g_S = g_n = (-)3.82$$

if the odd particle is a neutron. Since by assumption only the unpaired particle contributes to the nuclear spin I , the result must follow that $S = \frac{1}{2}$ or $S = -\frac{1}{2}$, so that $L = I + S$ must have either the value $L = I + \frac{1}{2}$ or $L = I - \frac{1}{2}$. Further, the parity of the state of the nucleus must be either odd or even, but cannot be both, so the orbital angular momentum ℓ of the unpaired nucleon must be either $\ell = I + \frac{1}{2}$ or $\ell = I - \frac{1}{2}$. With these assumptions it is possible to derive the Schmidt equations³⁷ in the following forms:

37. Fermi, E. Nuclear Physics. Chicago: The University of Chicago Press, 1950, pp. 19-21.

$$\mu_I = I + 2.29 \quad \text{odd proton, } I = \ell + \frac{1}{2},$$

$$\mu_I = \frac{I^2 - 1.29I}{I+1} \quad \text{odd proton, } I = \ell - \frac{1}{2},$$

$$\mu_I = -1.91 \quad \text{odd neutron, } I = \ell + \frac{1}{2}, \text{ and}$$

$$\mu_I = \frac{1.91I}{I+1} \quad \text{odd neutron, } I = \ell - \frac{1}{2}.$$

Since the assumptions require that $I = \ell \pm \frac{1}{2}$, the Schmidt equations thus will yield two values for the predicted μ_I of any particular nuclide. It is found in practice that nuclear magnetic moments observed to date will, with only a few exceptions, lie between the two values yielded by these equations. Those observed values which are exceptions fall outside of these limits only by comparatively small amounts.

is the one variable in a matrix. When a matrix is given, the matrix is usually written in the form $A = [a_{ij}]$, where i and j are subscripts. The value of a_{ij} is the element in the i th row and j th column. For example, if $A = [a_{ij}]$ is a 3×3 matrix, then a_{11} is the element in the first row and first column, a_{12} is the element in the first row and second column, and so on. The matrix A is said to be symmetric if $a_{ij} = a_{ji}$ for all i and j . The matrix A is said to be diagonal if $a_{ij} = 0$ for all $i \neq j$. The matrix A is said to be the identity matrix if $a_{ij} = 1$ for $i = j$ and $a_{ij} = 0$ for $i \neq j$. The matrix A is said to be the zero matrix if $a_{ij} = 0$ for all i and j .

27. Find the inverse of the matrix $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$.
 Answer: $A^{-1} = \frac{1}{-2} \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix} = \begin{bmatrix} -2 & 1 \\ 1.5 & -0.5 \end{bmatrix}$

old matrix $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$	$\mu_1 = 1 + 2i$
old matrix $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$	$\mu_1 = \frac{1 - 1.29i}{1 + i}$
old matrix $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$	$\mu_1 = -1.29i$
old matrix $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$	$\mu_1 = \frac{1.91i}{1 + i}$

When the eigenvalues are complex, the eigenvectors are also complex. If $\mu_1 = a + bi$ is an eigenvalue of a matrix A , then $\mu_2 = a - bi$ is also an eigenvalue of A . The eigenvectors corresponding to μ_1 and μ_2 are complex conjugates of each other. The matrix A is said to be normal if $AA^H = A^H A$, where A^H is the conjugate transpose of A . The matrix A is said to be Hermitian if $A = A^H$. The matrix A is said to be skew-Hermitian if $A = -A^H$. The matrix A is said to be unitary if $A^{-1} = A^H$. The matrix A is said to be orthogonal if $A^{-1} = A^T$, where A^T is the transpose of A .

Unfortunately, the difference between the two Schmidt limits extends over ranges covering differences from two to four nuclear magnetons. This lack of preciseness seriously limits the usefulness of the theory.

By adopting some of the results contained in the theory of closed shells in nuclei as proposed by Mayer³⁶ and others it is possible to limit the choice of ℓ for the odd particle to only one value. By this theory it is found that $\ell = 2$ for both Cl^{35} and Cl^{37} ; for In^{113} and In^{115} , $\ell = 4$.

The two values of the nuclear magnetic moments predicted by the Schmidt equations for Cl^{35} and Cl^{37} , for In^{113} and In^{115} , and for N^{14} together with the observed values are listed in Table 6. The values obtained by using the nuclear shell theory results for ℓ with the appropriate Schmidt equation to obtain a single value for each μ_I are indicated by asterisks.

The examples in Table 6 are too limited in number to indicate the proper degree of merit which should be attached to the above method of attempting to predict values of μ_I . That this procedure leaves much to be desired would be seen, however, if one were to consider the cases which should be predicted most accurately. These cases consist of nuclides with an even number of neutrons and one more proton than the number required to complete the filling of a shell. It would seem in these cases that fairly close agreement should be obtained between theory and experiment, for the filled "core" of protons should be spherically symmetric and the magnetic moment should be due only to the one additional proton. But even

University of Illinois, Urbana-Champaign, Urbana, Illinois 61801

of the Group.

Section 7701(b)(1) of the Internal Revenue Code provides that the estate of a decedent is the estate of the decedent at the time of his death. The estate of a decedent is the estate of the decedent at the time of his death.

© 1998 by the American Psychological Association or one of its allied publishers. This article is intended solely for the personal use of the individual user and is not to be disseminated broadly.

© 1997 by The American Psychological Association, 0893-3200/97/\$12.00 DOI: 10.1037/0893-3200.11.4.475

possible to limit the number of l for the odd variables to only one value. By this theory it is found that $l = 2$ for both l_1 and l_2 ; the first and last $l = 1$.

For the values of the different constants mentioned by the
different equations for U_1 , U_2 , U_3 , U_4 , U_5 , and for U_6
together with the observed values are listed in table 6. The values
obtained by using the constant shell theory results for U_1 with the
improvements U_2 to U_6 are also given in table 6.

The example in Table 2 was not included in the analysis. The number of tests which should be retained in the analysis of the data is given by the number of tests which are retained in the analysis of the data. The number of tests which are retained in the analysis of the data is given by the number of tests which are retained in the analysis of the data.

Table 6.

Predictions of μ_I From Schmidt Equations and Shell Theory

Nuclide	Spin	ℓ (Shell Theory)	μ_I (Schmidt Model)		μ_I (Observed)	Diff. (S.M. - obs.)
			Lower	Higher		
$^{170}\text{I}^{35}$	3/2	2	0.126*	3.79	0.820896	-0.695
$^{17}\text{Cl}^{37}$	3/2	2	0.126*	3.79	0.683722	-0.458
$^{49}\text{In}^{113}$	9/2	4	2.60	6.79*	5.49507	+1.29
$^{49}\text{In}^{115}$	9/2	4	2.60	6.79*	5.50945	+1.28
$^{7}\text{N}^{14}$	1	(- - - Not Applicable - - -)			0.40361	

* indicates value obtained using ℓ from nuclear shell theory.

in these cases the predictions are found to be no more accurate than those in many cases less favorable.³⁶

In^{113} and In^{115} , two of the isotopes represented in Table 6, belong to a class of isotopes having identical spins but having $N = A - Z$ differing by two, where N is the number of neutrons in the nuclide. Frequently, pairs or triplets of isotopes of this type have almost identical magnetic moments. Examples of these are Ag^{107} and Ag^{109} ; Cs^{133} , Cs^{135} , and Cs^{137} ; Tl^{203} and Tl^{205} ; and In^{113} and In^{115} . This indicates that the protons, which according to Schmidt theory are responsible for the magnetic moments of all of these even-neutron nuclei, do not have their arrangement very greatly disturbed when two more neutrons are added to the nucleus. In^{113} and In^{115} also have very similar quadrupole moments and isomeric levels.³⁶

(b) Relationships Between μ_z and Q

For nuclei containing closed proton shells plus or minus one, the nuclear shell model not only predicts the sign of the quadrupole moment, but if the state of this odd proton can be ascertained from the nuclear spin and the magnetic moment, it is possible to arrive at a fairly good value for the quadrupole moment.³⁸ The nuclear shell

38. Townes, C.H., Foley, H.V., and Low, W. Phys. Rev. 76, 1415 (1949).

model also leads to other less quantitative conclusions concerning quadrupole moments.

Townes and his collaborators³⁸ have proposed a simple model of the nucleus based on nuclear shell considerations which leads to the proper behavior of known quadrupole moments but leaves the predictions of some of their magnitudes appreciably in error. This model is characterized by the following:

1. Neutrons and protons fit into single particle levels in a scheme similar to those proposed for correlating spins, thus producing what may be called proton and neutron shells.

2. Proton and neutron shells tend to be oriented or polarized to allow maximum overlap between proton and neutron distributions.

This model is stated by its originators to lead to the following conclusions:

- A. For an odd-proton nucleus, the quadrupole moment is primarily dependent on the number of protons P and can be written $Q_{p \text{ odd}} = Q_p(P)$, in which Q_p is always positive immediately before,

For small deviations around the mean, the distribution is Gaussian. The Gaussian model is not only useful for the study of the distribution of the mean, but in the case of the variance, it is also useful for the study of the distribution of the variance. The Gaussian model is also useful for the study of the distribution of the variance.

For small deviations around the mean, the distribution is Gaussian. The Gaussian model is not only useful for the study of the distribution of the mean, but in the case of the variance, it is also useful for the study of the distribution of the variance. The Gaussian model is also useful for the study of the distribution of the variance.

For small deviations around the mean, the distribution is Gaussian. The Gaussian model is not only useful for the study of the distribution of the mean, but in the case of the variance, it is also useful for the study of the distribution of the variance. The Gaussian model is also useful for the study of the distribution of the variance.

For small deviations around the mean, the distribution is Gaussian. The Gaussian model is not only useful for the study of the distribution of the mean, but in the case of the variance, it is also useful for the study of the distribution of the variance. The Gaussian model is also useful for the study of the distribution of the variance.

For small deviations around the mean, the distribution is Gaussian. The Gaussian model is not only useful for the study of the distribution of the mean, but in the case of the variance, it is also useful for the study of the distribution of the variance. The Gaussian model is also useful for the study of the distribution of the variance.

For small deviations around the mean, the distribution is Gaussian. The Gaussian model is not only useful for the study of the distribution of the mean, but in the case of the variance, it is also useful for the study of the distribution of the variance. The Gaussian model is also useful for the study of the distribution of the variance.

For small deviations around the mean, the distribution is Gaussian. The Gaussian model is not only useful for the study of the distribution of the mean, but in the case of the variance, it is also useful for the study of the distribution of the variance. The Gaussian model is also useful for the study of the distribution of the variance.

and always negative immediately after a shell is filled.

B. For an odd-neutron nucleus, the magnitude of the quadrupole moment depends on the number of protons, but its sign is determined by the number of neutrons, N , being given by $(Q_p(N)/[Q_n(N)])$. In this expression $Q_n(N)$ is the electric quadrupole moment which would be produced if they were protons. $Q_n(N)$ is very nearly the same function as $Q_p(N)$.

C. For odd-odd nuclei, estimation of quadrupole moments is more complex and depends on the way in which the angular momenta of the odd neutron and odd proton add. If these moments are essentially parallel, the quadrupole moment should be of the same sign and approximately the same magnitude as for a similar odd-proton nucleus. If the neutron and proton angular momenta are not essentially parallel, the quadrupole moment magnitude should be considerably reduced.

Generally it may be said that the nuclear shell model does not give correct values of Q . This is in contrast to the nuclear magnetic moments, for which an appropriate admixture of states of one nucleon can account for the magnetic moment of the nucleus, although this procedure does not appear very plausible by even the presently known facts about the nucleus. Large quadrupole moments demand an appreciable contribution from the protons which, according to the shell model, are in closed shells. For this contribution to be realized, it appears that angular momentum would have to be shared between the protons of the incomplete and of the filled shells. This polarization and the large asymmetry of distribution of nucleons is not entirely compatible with the single-particle central field concept

which forms the foundation of shell theory.³⁸

Hence it may be said that there is no well understood relationship between μ_I^2 and Q . Both are dependent upon spin, and both are functions of the number of protons and neutrons in a nucleus. To date no theory has been successful in establishing any important relationship between these two quantities.

The present study is directed toward the determination of the relationship between μ_I^2 and Q for the various isotopes of the elements. The study is directed toward the determination of the relationship between μ_I^2 and Q for the various isotopes of the elements. The study is directed toward the determination of the relationship between μ_I^2 and Q for the various isotopes of the elements.

The study is directed toward the determination of the relationship between μ_I^2 and Q for the various isotopes of the elements. The study is directed toward the determination of the relationship between μ_I^2 and Q for the various isotopes of the elements. The study is directed toward the determination of the relationship between μ_I^2 and Q for the various isotopes of the elements.

[illegible]

ACKNOWLEDGMENTS

The work reported in this thesis was carried out while the writer was undergoing post graduate training in Radiological Defense Engineering at the Ohio State University under a program sponsored by the Armed Forces Special Weapons Project and supervised by the United States Naval Post Graduate School at Monterey, California. A part of the equipment used in this work was supplied under a research contract between the Ohio State University Research Foundation and the Flight Research Laboratory of the U.S. Air Force Wright-Patterson Air Development Center.

The writer wishes to express his sincerest appreciation for the extensive advice and encouragement given him in the above work by his adviser, Professor Dudley Williams, and to Dr. Yu Ting for his extremely valuable assistance in all phases of the work covered in this thesis.

APPENDIX

The work reported in this thesis was carried out while the author was a member of the staff of the Department of Zoology, University of California, Berkeley, during the summer of 1934. The work was carried out under the supervision of Professor J. B. Clark, and the results are presented in this thesis. The work was supported in part by a grant from the National Science Foundation, and in part by a grant from the University of California.

The author wishes to express his appreciation to Professor J. B. Clark for his generous hospitality and for the opportunity to work in his laboratory. He also wishes to express his appreciation to the members of the staff of the Department of Zoology, University of California, Berkeley, for their helpful criticism and suggestions. The author also wishes to express his appreciation to the members of the staff of the Department of Zoology, University of California, Berkeley, for their helpful criticism and suggestions.

Thesis
B51

19389

Biard

Nuclear magnetic mo-
ments of nitrogen¹⁴,
chlorine³⁷, and indium
¹¹⁵.

Thesis
B51

19389

Biard

Nuclear magnetic moments of
nitrogen¹⁴, chlorine³⁷, and
indium¹¹⁵.

Library

U. S. Naval Postgraduate School

Monterey, California

thesB51

Nuclear magnetic moments of nitrogen(14)



3 2768 002 13437 1

DUDLEY KNOX LIBRARY